

CEE 110 Probability and Statistics for Engineers and Scientists  
Midterm


Good luck!! Please show ALL work.

Problem 1:

A pound is trying desperately to tame many dogs for adoption. Both giving treats and playing with the dogs are factors contributing to how friendly a dog will be after a week-long training program. They do an experiment to find out how best to efficiently tame the dogs. Considering the following data on average dog friendliness in the four different groups at the end of the program, please calculate main effects for each factor, identify whether the factors are additive or interaction, and please quantify any interaction, if any. (Solve for the mean you'd expect for dogs treated with both treats and playtime if the factors were additive.)

	No treats	With treats	Row averages	Main row effects
No playtime	30	70	50	-11.25
With playtime	50	95	72.5	+11.25
Column averages	40	82.5	61.25	
Main column effects	-21.25	+21.25		

Please draw the graph used to visualize interaction (one factor along x axis and lines drawn for each case for the other factor).



There are not parallel so we can see that the factors have interaction.

$$Y_{ij} = \mu_{ij} - (\alpha_i + \beta_j + \theta_{ij})$$

$$Y_{\text{no playtime, no treats}} = 30 - (\alpha_1 + \beta_1 + \theta_{11}) = 12.5$$

$$Y_{\text{no playtime, with treats}} = 70 - (\alpha_1 + \beta_2 + \theta_{12}) = 12.5$$

$$Y_{\text{with playtime, no treats}} = 50 - (\alpha_2 + \beta_1 + \theta_{21}) = 12.5$$

$$Y_{\text{with playtime, with treats}} = 95 - (\alpha_2 + \beta_2 + \theta_{22}) = 12.5$$

mean for dog w/ treats and w/ playtime if factors were additive

$$\mu_{ij} = \mu_{..} + \alpha_{\text{with playtime}} + \beta_{\text{with treats}} = 61.25 + 11.25 + 21.25 = 93.75$$

Problem 2: Short answer:

A) More and more fuses are selected until four defective fuses are found. The number of fuses tested up to and including the fourth defective fuse is recorded.

$$S_X = (4, 5, 6, \dots, \infty)$$

B) Three students from this class are selected to each rate a new movie on a scale of 1-4 stars. The sample space contains how many possible outcomes?


$$S_1 = \{1, 2, 3, 4\} \quad S_2 = \{1, 2, 3, 4\} \quad S_3 = \{1, 2, 3, 4\}$$

$$4 \cdot 4 \cdot 4 = \boxed{64 \text{ possible outcomes}}$$

C) What are the odds of two dice summing to ten?

$$P = \frac{N(E)}{N}$$

$$\frac{3}{36} = \boxed{\frac{1}{12}} \text{ probability}$$



D) What is the probability of rolling at least one double "1's" in sixteen rolls of two dice?

$$1 - \left(\frac{35}{36}\right)^{16} = 1 - .637 = \boxed{0.36284}$$

100

Problem 3:

A leadership committee of four students is to be formed from ten students.

- a) How many different committees are possible?
- b) The ten students are five biology majors, three chemists, and two physicists. How many committees would have two biology majors, one chem major, and one physics major?
- c) If all committees were equally likely to form, what is the likelihood of a committee as outlined in part b.

a)  $\binom{10}{4} = \boxed{210}$  committees possible

$$\frac{10!}{4!6!}$$

b)  $\binom{5}{2} \binom{3}{1} \binom{2}{1} = (10)(3)(2) = \boxed{60}$  committees

$$\frac{5!}{2!3!} \frac{3!}{2!1!} \frac{2!}{1!1!}$$

c)  $\frac{60}{210} = \boxed{0.2857}$


$$\frac{\binom{5}{2} \binom{3}{1} \binom{2}{1}}{\binom{10}{4}}$$

Problem 4:

Eighty-five percent of all students have smart phones while 15% have flip phones. Of those with smart phones, 60% check email with high frequency (every hour or two), 30% check email with middle frequency (three or four times a day) and 10% check it with low frequency (once or twice per day). Of those using flip phones, 20%, 40%, and 40% have incomes in the high, middle and low categories, respectively.

Please draw tree diagram

- a) What is the probability that a randomly chosen student only checks email once or twice a day? = low  
b) Given a low frequency of email checking, what is the probability that s/he has a smart phone?



$$a) 0.85(0.10) + 0.15(0.40) = 0.085 + 0.06 = \boxed{0.145}$$

$$b) P(\text{smartphone} | \text{low freq}) = \frac{P(\text{low freq} \cap \text{smartphone})}{P(\text{low freq})}$$

$$= \frac{.85(.10)}{.145} = \boxed{.58621}$$

$$P(A \cap B) = P(A) P(B|A)$$
$$P(B|A) = \frac{P(A \cap B)}{P(A)}$$

Problem 5:

On average, 2.4 spotted owls land in a certain grove of Redwoods per hour.

- What is the probability that four or less birds will fly by in an hour?
- What is the probability that two, three, or four birds will fly by in an hour?
- What is the probability that at least two birds will fly by in the hour?

$$\lambda = 2.4$$

$$a) P(X \leq 4) = \boxed{0.904} \text{ taken from table}$$

$$b) P(1 < X \leq 4) = P(X \leq 4) - P(X \leq 1) = 0.904 - 0.308 = \boxed{0.596}$$

$$c) P(X \geq 2) = 1 - P(X \leq 1) = 1 - 0.308 = \boxed{0.692}$$

Problem 6:

A habitat contains 20 snow leopards, of which five are fitted with a radio collar. Six snow leopards are captured at a later time, and let  $X$  be the number of collared snow leopards in the group (assuming the collar does not affect the probability of being captured).

a) Is  $X$ :

- a. Binomial
- b. Hypergeometric
- c. Negative binomial
- d. Poisson

b) What is the probability that  $X = 2$ ?

c) What is the expected value of  $X$ ?

a) hypergeometric,  $n = 6$ ,  $M_1 = 5$ ,  $M_2 = 20 - 5 = 15$

$$b) P(X=x) = \frac{\binom{M_1}{x} \binom{M_2}{n-x}}{\binom{M_1+M_2}{n}} = \frac{\binom{5}{2} \binom{15}{4}}{\binom{20}{6}} = \frac{\left(\frac{5!}{2!3!}\right) \left(\frac{15!}{11!4!}\right)}{\left(\frac{20!}{14!6!}\right)} = \boxed{0.3522}$$

$$c) E(X) = M_x = n \frac{M_1}{N} = 6 \left(\frac{5}{20}\right) = \boxed{1.5}$$

This page is left blank for your use.


Problem 7:

Adult frogs of a certain species are being studied for effects due to climate change. Their length is normally distributed with mean length of 4.2 cm and standard deviation 0.3 cm. An adult frog is considered in a health range if its length is between 3.5 and 4.6 cm.

- a) What is the probability that a randomly captured frog will be in the healthy range?
- b) What is the probability that of four randomly and independently captured frogs, two are acceptable?

Please draw a normal curve and mark off the area you are solving for.

$\mu = 4.2 \text{ cm}$      $\sigma = 0.3 \text{ cm}$



$$\begin{aligned}
 a) \ P(3.5 \leq X \leq 4.6) & \Rightarrow \Phi\left(\frac{4.6 - 4.2}{0.3}\right) - \Phi\left(\frac{3.5 - 4.2}{0.3}\right) \\
 & = \Phi\left(\frac{0.4}{0.3}\right) - \Phi\left(\frac{-0.7}{0.3}\right) \\
 & = \Phi(1.33) - \Phi(-2.33) \\
 & = \Phi(1.33) - (1 - \Phi(2.33)) \\
 & = 0.9082 - (1 - .9901) = \boxed{0.8983}
 \end{aligned}$$

$$b) \ \binom{4}{2} (.8983)^2 (1 - .8983)^2 = \frac{4!}{2!2!} (.8983^2) (.1017^2) = \boxed{0.0501}$$

that two will be acceptable