

NAME: Mike Wilson !!
003 506374

CS170A – Mathematical Models and Methods for Computer Science
Spring 2007
D.S. Parker

Midterm Examination
OPEN BOOK, OPEN NOTES
Thursday, May 10, 2007, 4:00pm-5:50pm

Do not cheat.

<i>Problem</i>	<i>Points</i>
1	19/25
2	25/25
3	25/25
4	25/25
<i>Total</i>	94/100

1. Everything

For each of the following matrices, determine (yes/no) whether the matrix is: unitary, hermitian, invertible, positive definite. (Recall A is positive definite if it is Hermitian and all of its eigenvalues are positive real values.) Assume that $i = \sqrt{-1}$.

matrix	unitary?	hermitian?	invertible?	positive definite?
$\det = 0-2 = -2$ $\begin{pmatrix} 0 & 1 \\ 2 & 3 \end{pmatrix}$	True <input type="checkbox"/> False <input checked="" type="checkbox"/>	True <input type="checkbox"/> False <input checked="" type="checkbox"/>	True <input checked="" type="checkbox"/> False <input type="checkbox"/>	True <input type="checkbox"/> False <input checked="" type="checkbox"/>
$\det = -1$ $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$	True <input checked="" type="checkbox"/> False <input checked="" type="checkbox"/>	True <input checked="" type="checkbox"/> False <input type="checkbox"/>	True <input checked="" type="checkbox"/> False <input type="checkbox"/>	True <input type="checkbox"/> False <input checked="" type="checkbox"/>
$\det = 2$ $\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$	True <input type="checkbox"/> False <input checked="" type="checkbox"/>	True <input checked="" type="checkbox"/> False <input type="checkbox"/>	True <input checked="" type="checkbox"/> False <input type="checkbox"/>	True <input type="checkbox"/> False <input checked="" type="checkbox"/>
$\det = 3-2 = 1$ $\begin{pmatrix} 1 & -1 \\ -2 & 3 \end{pmatrix}$	True <input checked="" type="checkbox"/> False <input checked="" type="checkbox"/>	True <input type="checkbox"/> False <input checked="" type="checkbox"/>	True <input checked="" type="checkbox"/> False <input type="checkbox"/>	True <input type="checkbox"/> False <input checked="" type="checkbox"/>

For each of the following equations, mark whether the equation is True (valid for all specified matrices A) or False. Assume that A' denotes the transpose of A , A^H denotes the hermitian transpose of A , and $i = \sqrt{-1}$.

For full credit, if you mark it True, you must explain how you derived this. If you mark it False, you must give a counterexample (values for these sets for which the equation does not hold).

(a) True False

If $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ then the eigenvalues of A are $\lambda = ((a+d) \pm \sqrt{(a+d)^2 - 4(ad-bc)})/2$

Explanation: This is the soln to the eqn $\det(\lambda I - A) = 0$
see previous page

(b) True False

If $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ then $\text{tr}(A) = a+d$ (= the sum of the diagonal elements of A) is the same as the sum of the eigenvalues of A .

Explanation: This true if A has real eigenvalues.
But in $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ they are not.

(c) True False

Given a rotation matrix $Q = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix}$ with rotation angle θ , then Q^n is a rotation matrix with rotation angle $n\theta$.

Explanation: if $n=2$, $Q^2 = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} \cos^2 \theta + \sin^2 \theta & -\sin \theta \cos \theta - \sin \theta \cos \theta \\ 2 \sin \theta \cos \theta & -\sin^2 \theta + \cos^2 \theta \end{pmatrix} = \begin{pmatrix} 1 & -2 \sin \theta \cos \theta \\ 2 \sin \theta \cos \theta & \cos^2 \theta - \sin^2 \theta \end{pmatrix}$

(d) True False

If $A = \text{cov}(X)$ is a covariance matrix, then its eigenvalues are real and nonnegative.

Explanation: Since $\text{cov}(X)$ is of the form $M^H M$ for some $[M]_{n \times m}$, then if $M = USV^H$, $M^H M = VS^H U^H U S V^H = VS^2 V^H$

2. SVD

Assume that X is a real matrix, and that A is the covariance matrix $\text{cov}(X)$, so

$$A = \text{cov}(X) = \frac{1}{n-1} (X - M)' (X - M)$$

where M is the matrix whose columns are filled with means of corresponding columns in X .

(a) Show that A must have an SVD of the form

let $X - M = B$ $A = Q D Q'$

where D is a diagonal matrix of nonnegative real values and Q is orthogonal.

By SVD, $B = U S V'$, where S is diagonal of nonnegative real values, U, V are orthogonal.

then $B' B = V S' U' U S V' = V S^2 V' \sim Q D Q'$ with these properties $U' U = I$ since its orthogonal

(b) Assuming again that $A = Q D Q'$, where D and Q are as in part (a), show that A is positive definite.

D contains the eigenvalues of A , and all are positive real values.

Since all matrices of the form $M' M$ are symmetric (assuming real values)

and $A \sim (X - M)' (X - M)$ with real values, $A = A^T \Rightarrow A$ is Hermitian.

A is Hermitian and all eigenvalues are positive real $\Rightarrow A$ is pos. definite.

(c) Assuming again that $A = Q D Q'$, and using the fact that $\det(XYZ) = \det(X) \det(Y) \det(Z)$ for square matrices X, Y, Z , give a formula for $\det(A)$ in terms of $\det(D)$.

$$\det(A) = \det(Q) \det(D) \det(Q')$$

since Q and Q' are orthogonal, $\det(Q) = \det(Q') = 1$

$$\det(A) = 1 \cdot \det(D) \cdot 1 = \det(D)$$

(d) Assuming that $A = \text{cov}(X)$, is A Hermitian? If yes, show why; if not, give a counterexample.

Yes, because it is the product of a matrix with its conj. transp., which is a symmetric matrix.

Since A is symmetric, it must be Hermitian since $A = A'$.

(e) Assuming again that $A = Q D Q'$, what is the SVD of A^{-1} ?

$$A^{-1} = (Q D Q')^{-1}$$

$$= (Q')^{-1} D^{-1} Q^{-1}$$

$$= Q D^{-1} Q' \quad \text{since } Q, Q' \text{ are orthogonal, and } Q' = Q^{-1}$$

3. Least Squares

Gauss became famous because he was able to fit an ellipse through the observed positions of the asteroid Ceres using least squares. Your job here is to show how he obtained this fit.

Suppose you are given a table of n pairs of real (x, y) values

x	y
x_1	y_1
\vdots	\vdots
x_n	y_n

and you are asked to fit the line

$$ax^2 + by^2 = 1$$

through this data, finding real coefficients a and b that minimize squared error.

- (a) Express this as a linear problem. That is, give a matrix A whose entries are functions of $\{x_i\}$ and $\{y_i\}$ for which a and b can be obtained by least squares on $A \begin{pmatrix} a \\ b \end{pmatrix} = \vec{1}$, where $\vec{1}$ is the $n \times 1$ vector whose entries are all 1.

$$A = \begin{pmatrix} x_1^2 & y_1^2 \\ \vdots & \vdots \\ x_n^2 & y_n^2 \end{pmatrix} \Rightarrow \begin{pmatrix} a \\ b \end{pmatrix} = A \begin{matrix} \vec{1} \\ \uparrow \\ \text{fancy} \end{matrix}$$

- (b) Give an explicit formula for $A'A$ (in terms of $\{x_i\}$ and $\{y_i\}$).

$$A'A = \begin{pmatrix} \sum x_i^4 & \sum x_i^2 y_i^2 \\ \sum x_i^2 y_i^2 & \sum y_i^4 \end{pmatrix}$$

(1+2)(2+4)

- (c) Give an explicit formula for a (in terms of $\{x_i\}$ and $\{y_i\}$).

$$a = (A'A)^{-1} A' \vec{1}$$

$$A' \vec{1} = \begin{pmatrix} x_1^2 & \dots & x_n^2 \\ y_1^2 & \dots & y_n^2 \end{pmatrix} \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} = \begin{pmatrix} \sum x_i^2 \\ \sum y_i^2 \end{pmatrix}$$

$$(A'A)^{-1} A' = \frac{1}{\sum x_i^4 \sum y_i^4 - 2 \sum x_i^2 y_i^2} \begin{pmatrix} \sum y_i^4 & -\sum x_i^2 y_i^2 \\ -\sum x_i^2 y_i^2 & \sum x_i^4 \end{pmatrix}$$

$$a = \frac{\sum y_i^4 \sum x_i^2 - \sum x_i^2 y_i^2 \sum y_i^2}{\sum x_i^4 \sum y_i^4 - 2 \sum x_i^2 y_i^2}$$

OK

4. PCA

Consider the following list of polyhedra and their properties:

Polyhedron	Faces	Vertices	Edges
tetrahedron	4	4	6
octahedron	8	6	12
cube	6	8	12
dodecahedron	12	20	30
icosahedron	20	12	30

With Matlab we produce following session:

```
>> X = [
    4    4    6 ;
    8    6   12 ;
    6    8   12 ;
   12   20   30 ;
   20   12   30 ;
];

>> cov(X)
ans =
    40    23    63
    23    40    63
    63    63   126

>> [U,S,V] = svd(cov(X))
U =
   -0.4082    0.7071   -0.5774
   -0.4082   -0.7071   -0.5774
   -0.8165   -0.0000    0.5774
S =
  189.0000     0     0
     0   17.0000     0
     0     0    0.0000
V =
   -0.4082    0.7071   -0.5774
   -0.4082   -0.7071   -0.5774
   -0.8165   -0.0000    0.5774
>>
```

- (a) Identify the first two principal components of the dataset X .

$$U(:,1) \text{ and } U(:,2) = (.7071; -0.7071; 0)$$

$$\hookrightarrow = (-.4082; -.4082; -.8165)$$

- (b) Explain how to project the dataset X onto these first two components.

The dot product of each with X . ex. $\begin{bmatrix} 4 & 4 & 6 \\ 8 & 6 & 12 \\ 6 & 8 & 12 \\ 12 & 20 & 30 \\ 20 & 12 & 30 \end{bmatrix} \begin{bmatrix} -.4082 \\ -.4082 \\ -.8165 \end{bmatrix} = \begin{bmatrix} \dots \\ \dots \\ \dots \\ \dots \\ \dots \end{bmatrix}$

- (c) Suppose we want to find a linear relationship between the number of Faces, Vertices, and Edges. That is, we want to find real values a , b , c such that

$$\text{Faces} = a \cdot \text{Vertices} + b \cdot \text{Edges} + c$$

Explain how these equations could be solved using least squares, by converting them to matrix form and explaining what matrix operations you could use to find a , b , and c .

Use $A = \begin{pmatrix} V_1 & E_1 & 1 \\ \vdots & \vdots & \vdots \\ V_n & E_n & 1 \end{pmatrix}$ then $A \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} F_1 \\ \vdots \\ F_n \end{pmatrix} \Rightarrow \begin{pmatrix} a \\ b \\ c \end{pmatrix} = (AA^T)^{-1} A^T \begin{pmatrix} F_1 \\ \vdots \\ F_n \end{pmatrix}$

- (d) The last singular value is zero; what does this mean about the value of c ? the dataset X

That means the third (edges) follows from the first

two, faces & vertices.

This implies that you only need
the first 2 PCs to reconstruct the dataset.