


Full Name ~~XXXXXXXXXX~~

Yanli Liu	1A T 9:00A-9:50A MS 6229 1B R 9:00A-9:50A Boelter 5419
Nicholas Boschert	1C T 9:00A-9:50A Boelter 5280 1D R 9:00A-9:50A Boelter 5280
Gyu Eun Lee	1E T 9:00A-9:50A Boelter 5272 1F R 9:00A-9:50A Boelter 5272

Section	I	E
---------	---	---

- Do not open this exam packet until you are told that you may begin.
- Turn off all electronic devices and and put away all items except for a pen/pencil and an eraser.
- Remove hats and sunglasses.
- No phones, calculators, smart-watches or electronic devices of any kind allowed for any reason, including checking the time.
- If you have a question, raise your hand and one of the proctors will come to you. We will not answer any mathematical questions except possibly to clarify the wording of a problem.
- Quit working and close this packet when you are told to stop.



Page:	1	2	3	4	5	6	7	8	Total
Points:	6	4	8	7	6	8	5	6	50
Score:	6	4	8	4	6	8	5	6	47

Part I: Multiple choice. Please write your answers (A, B, C, ...) in the boxes on the right.

1. (3 points) Determine values of the constants a and b that make the differential equation exact.

$$(3x^2y - bx^5y^3) dx + (ax^3 + x^6y^2) dy = 0 \quad (1)$$

- A. $a = 2, b = -1$
- B. $a = 1, b = -2$
- C. $a = 2, b = 1$
- D. $a = -1, b = 2$
- E. None of the above.

$$P_y = 3x^2 - bx^5(3y^2)$$

$$Q_x = a_1 = a(3x^2) + 6x^5y^2$$

$$a = 1, b = -2$$


B

2. (3 points) Consider the initial value problem

$$y' = (y^2 - 1) \sin^2(ty), \quad y(0) = 2.$$


Which of the following is true?

- A. $-1 < y(t) < 1$ for all t for which y is defined.
- B. $y(t) < \sin^2(t)$ for all t for which y is defined.
- C. $y(t) > 1$ for all t for which y is defined.
- D. $y(t) > t^2 - 1$ for all t for which y is defined.
- E. None of the above.



C

3. (4 points) The slope field below corresponds to which differential equation?



(A) $y' = \frac{t^2}{1-y^2}$

b/c even with respect to both t and y

B. $y' = t^2$

C. $y' = 4y$

D. $y' = ty$

E. $y' = \frac{y^2}{2t+1}$

A

4. (4 points) Determine if the equation

$$\left(\frac{y}{x} + 8x\right) dx + (\ln x - 3) dy = 0 \quad (2)$$

is exact (you may assume that we are working in a rectangle R in the plane such that $x > 0$ for all (x, y) in R). If it is exact, find the solution.

A. $F(x, y) = y \ln x + 4x^2 - 3y$

B. $F(x, y) = -\frac{y}{x^2} + \frac{1}{x} + 8$

C. $-\frac{y}{x^2} + \frac{1}{x} + 8 = C$

D. $y \ln x + 4x^2 - 3y = C$

E. The equation is not exact.

$P_y = \frac{1}{x} \quad Q_x = \frac{1}{x}$

$\frac{\partial F}{\partial x} = \frac{y}{x} + 8x \quad \frac{\partial F}{\partial y} = \ln x - 3$

$\frac{\partial F}{\partial x} = \frac{y}{x} + 8x \quad \frac{\partial F}{\partial y} = \ln x - 3$

$F = y \ln x + 4x^2 - 3y = C$

D

5. (4 points) Which of the following integrating factors is suitable for the differential equation

$$(x + 2) \sin y dx + x \cos y dy = 0? \quad (3)$$

A. $e^{\cos x}$

B. $\sin x$

C. xe^x

D. $1 + \frac{1}{x}$

E. None of the above.

$\frac{1}{Q} [P_y - Q_x]$

$= \left(\frac{1}{x \cos y}\right) ((x+2) \cos y - \cos y)$

$= \frac{x+1}{x} = 1 + \frac{1}{x}$

$\int (1 + \frac{1}{x}) dx = x + \ln x$

$e^{x+\ln x} = e^x e^{\ln x} = x e^x$

C


6. (4 points) The function $\mu(x, y) = \frac{1}{x^2 + y^2}$ is an integrating factor for the equation (4)
- $$(x^2 + y^2 - x)dx - y dy = 0.$$

Use this to solve the differential equation. You may assume that we are working in a rectangle R which does not contain the point $(0, 0)$.

- A. $F(x, y) = x - \arctan(x^2 + y^2)$
 B. $F(x, y) = x - \frac{1}{2} \ln(x^2 + y^2)$
 C. $x - \frac{1}{2} \ln(x^2 + y^2) = C$
 D. $x - \arctan(x^2 + y^2) = C$
 E. None of the above

$$\begin{aligned} \frac{\partial F}{\partial x} &= \frac{x^2 + y^2 - x}{x^2 + y^2} & \frac{\partial F}{\partial y} &= \frac{-y}{x^2 + y^2} \\ &= 1 - \frac{x}{x^2 + y^2} \end{aligned}$$

$$F = x - \frac{1}{2} \ln(x^2 + y^2) = C$$



7. (3 points) True or false: there exists a differential equation of the form $y' = f(t, y)$ such that f has continuous partial derivatives on a rectangle R containing $(0, 0)$ and such that (5)

$$y_1 = 2t \quad \text{and} \quad y_2 = 3t$$

are both solutions in R .

- A. False. The existence theorem forbids it.
 B. False. The uniqueness theorem forbids it.
 C. True. The existence theorem guarantees it.
 D. True. The uniqueness theorem guarantees it.



Part II: Free Response. Write up a full solution for each problem. A correct answer with an incomplete or incorrect solution will not receive full credit.

8. (6 points) Find the general solution of the linear equation

$$y' - 2y = 4t^3 e^{2t}.$$

Box your answer

1) homogeneous equation

$$\frac{dy}{dt} = 2y \Rightarrow \frac{dy}{y} = 2dt \Rightarrow \ln|y| = 2t + C \Rightarrow y = Ae^{2t}$$

2) vary A, let $A = v(t)$

$$y = v(t)e^{2t}$$

3) Plug into original equation

$$y' - 2y = 4t^3 e^{2t} \Rightarrow y' = 2y + 4t^3 e^{2t}$$

$$2v(t)e^{2t} + v'(t)e^{2t} = \cancel{2v(t)e^{2t}} + 4t^3 e^{2t}$$

$$v'(t)e^{2t} = 4t^3 e^{2t}$$

$$v(t) = \int 4t^3 dt = t^4 + C$$

4) Put it back together

$$y = (t^4 + C)e^{2t}$$

9. (8 points) Consider two tanks, labeled tank A and tank B. Tank A contains 150 gal of solution in which is dissolved 12 lbs of salt. Tank B contains 300 gal of solution in which is dissolved 28 lbs of salt. Pure water flows into tank A at a rate of 4 gal/s. There is a drain at the bottom of tank A. The solution leaves tank A via this drain at a rate of 4 gal/s and flows immediately into tank B at the same rate. A drain at the bottom of tank B allows the solution to leave tank B at a rate of 1 gal/s. Set up, but do not evaluate, a system of differential equations involving the variables x (amount of salt in tank A), y (amount of salt in tank B), and t (time in seconds). Do not forget to include any initial conditions. Box your entire answer

A: at $t=0$: 150 gal
12 lb salt
in: 4 gal/s, 0 lb/gal
out: 4 gal/s x

B: at $t=0$: 300 gal
28 lb salt
in: 4 gal/s
out: 1 gal/s y

$$V(t) = 300 + 3t$$

A: rate of change = rate in - rate out

$$\begin{aligned} \frac{dx}{dt} &= \left(\frac{4 \text{ gal}}{s}\right)\left(\frac{0 \text{ lb}}{\text{gal}}\right) - \left(\frac{4 \text{ gal}}{s}\right)\left(\frac{x \text{ lb}}{150 \text{ gal}}\right) \\ &= \frac{-4x}{150} \Rightarrow \frac{dx}{dt} = \frac{-2x}{75} \end{aligned}$$

B: $\frac{dy}{dt} = \frac{2x}{75} - \left(\frac{1 \text{ gal}}{s}\right)\left(\frac{y \text{ lb}}{300+3t \text{ gal}}\right)$

$$\frac{dy}{dt} = \frac{2x}{75} - \frac{y}{300+3t}$$

$$\begin{aligned} \frac{dx}{dt} &= \frac{-2x}{75} \\ \frac{dy}{dt} &= \frac{2x}{75} - \frac{y}{300+3t} \\ x(0) &= 12, y(0) = 28 \end{aligned}$$

10. (5 points) The differential equation

$$y'' + 2t(y')^2 = 0$$

is an example of a second order differential equation. The change of variable $v = y'$ (so that $v' = y''$) turns this equation into a first-order separable equation. Using this change of variable, find the particular solution which satisfies

$$y(1) = \frac{\pi}{2}, \quad y'(0) = 1.$$

Box your answer

$$y'' + 2t(y')^2 = 0.$$

Let $v = y'$ and $v' = y''$. Then $v' + 2tv^2 = 0$.

$$\frac{dv}{dt} = -2tv^2 \Rightarrow$$

$$\frac{dv}{v^2} = -2t dt \Rightarrow$$

$$\frac{-1}{v} = -t^2 + C$$

$$\Rightarrow \frac{1}{v} = t^2 + C \Rightarrow v = \frac{1}{t^2 + C}$$

$$y'(0) = 1 \Rightarrow v(0) = 1 \Rightarrow 1 = \frac{1}{0 + C} \Rightarrow C = 1 \Rightarrow v = y' = \frac{1}{t^2 + 1}$$

$$y = \int \frac{1}{t^2 + 1} dt = \arctan(t) + D$$

$$y(1) = \frac{\pi}{2} \Rightarrow \frac{\pi}{2} = \arctan(1) + D \Rightarrow \frac{\pi}{2} = \frac{\pi}{4} + D \Rightarrow D = \frac{\pi}{4}$$

$$\Rightarrow \boxed{y = \arctan(t) + \frac{\pi}{4}}$$



11. (6 points) The differential equation

$$y^2 y' = -y^3 + 3e^{-t} \quad (6)$$

is an example of a Bernoulli equation with degree $n = -2$.

(a) The change of variables $z = y^3$ turns this equation into a first-order linear differential equation. Using this, write the linear equation above in the form $z' = a(t)z + f(t)$. (I will not be grading your work for this problem, just your answer). Box your answer.

$$z = y^3 \Rightarrow \frac{dz}{dy} = 3y^2$$

$$dz = 3y^2 dy$$

$$\frac{dz}{dt} = 3y^2 \frac{dy}{dt}$$

$$y^2 y' = -y^3 + 3e^{-t}$$

$$\frac{1}{3} \frac{dz}{dt} = -z + 3e^{-t}$$

$$\Rightarrow \boxed{z'(t) = -3z + 9e^{-t}}$$

(b) Solve the differential equation. Your answer should be in the form $y(t) = \dots$

1) hom eq

$$\frac{dz}{dt} = -3z \Rightarrow \frac{dz}{z} = -3dt \Rightarrow \ln|z| = -3t + C \Rightarrow z = Ae^{-3t}$$

2) let $A = v(t)$

$$z = v(t)e^{-3t}$$

3) Plug into original

$$z' = -3z + 9e^{-t}$$

$$-3v(t)e^{-3t} + v'(t)e^{-3t} = -3v(t)e^{-3t} + 9e^{-t}$$

~~$$v(t)e^{-3t} = 9e^{-t}$$~~

$$v'(t)e^{-3t} = 9e^{-t}$$

$$\Rightarrow v'(t) = 9e^{-t} e^{3t} = 9e^{2t}$$

$$v(t) = \frac{9}{2} e^{2t} + C$$

$$z(t) = \left(\frac{9}{2} e^{2t} + C\right) e^{-3t}$$

$$y = (z)^{1/3} \Rightarrow$$

$$\boxed{y(t) = \left[\left(\frac{9}{2} e^{2t} + C\right) e^{-3t}\right]^{1/3}}$$