

MT1 Physics 1B W16

Full Name (Printed) XIN LING

Full Name (Signature) Minaling


Student ID Number 704574468

Seat Number _____

Problem	Grade
1	08/30
2	14/30
3	19/30
Total	31 /90

41

- Do not peek at the exam until you are told to begin. You will have approximately 50 minutes to complete the exam.
- Don't spend too much time on any one problem. Solve 'easy' problems first. Go for partial credit!
- **HINT:** Focus on the concepts involved in the problem, the tools to be used, and the set-up. If you get these right, all that's left is algebra.
- **Have Fun!**



1) A pair of masses, M_1 and M_2 are joined by a spring of constant k and natural (unstretched) length L_0 .

- +5 • 1a) (5 points) Find the amount the spring is extended or compressed (Δx) as a function of the position of each mass (x_1 and x_2 , respectively). Make sure Δx is positive when the spring is extended and negative when it's compressed.

$$\Delta x = (x_2 - x_1) - L_0$$

- +3 • 1b) (5 points) Use Newton's laws to obtain differential equations (written in terms of x_1 and x_2) that describe the motion of each mass. Note that these are 'coupled' equations - they will each depend on both x_1 and x_2 - but not to worry, we'll address that later. Make sure the derivative term in each equation has the correct sign when the spring is extended and compressed!

~~F₁ = kΔx = m₁a₁~~ ~~positive because direction is opposite~~

$$m_1 \frac{d^2 x_1}{dt^2} = -k[(x_2 - x_1) - L_0] \quad x_1(t) = A \cos\left[\sqrt{\frac{k}{m_1}}(x_2 - x_1 - L_0)\right]$$

$$\frac{d^2 x_1}{dt^2} + \frac{k}{m_1}(x_2 - x_1 - L_0) = 0 \rightarrow -A \sin\left[\sqrt{\frac{k}{m_1}}(x_2 - x_1 - L_0)\right] + x_1$$

according to same principle:

$$\frac{d^2 x_2}{dt^2} + \frac{k}{m_2}(x_2 - x_1 - L_0) = 0$$

~~opposite direction~~

$$x_2(t) = A \sin\left[\sqrt{\frac{k}{m_2}}(x_2 - x_1 - L_0)\right] + x_2$$



- 1c) (5 points) Algebraically relate the derivative term in x_1 to the derivative term in x_2 . Use this result, along with your answer to part *a*, to obtain relationships between a derivative of Δx and each of the derivatives of x_1 and x_2 that appear in your answers to part *b*.

The image shows handwritten scribbles and a diagram. The scribbles are dense and cover a significant portion of the page. Below them is a diagram of a two-mass system. It consists of two masses, labeled m_1 and m_2 , connected by a spring. The masses are represented by horizontal bars. The spring is represented by a coiled line between the two bars. The masses are positioned on a horizontal surface, indicated by a line below them. The diagram is drawn in black ink.

- 1d) (10 points) Now it's time to put it all together - rewrite the differential equations from part *b* in terms of Δx . The results should look familiar. Find the solution for Δx as a function of time (make sure you evaluate, in terms of given information, any constants that are determined by the construction of the system).

- 1e) (5 points) Suppose we were to tie one end of the spring off to a wall, and the other end of the spring to a mass M . What value would M have to have in order for this new system to oscillate with the same period as the two-mass system we've been working on? This value, known as the *reduced mass*, is used to simplify the discussion of binary systems (diatomic molecules, for instance) in oscillation.

$$\omega_{damp} = \sqrt{\omega_0^2 - \left(\frac{b}{2m}\right)^2}$$

2) Recall that the amplitude of a driven mass-spring system is given by

$$A(\Omega) = \frac{F_0/m}{\sqrt{(\omega_0^2 - \Omega^2)^2 + \left(\frac{b\Omega}{m}\right)^2}}$$

- 2a) (10 points) Show that the following relationship holds true for a mass-spring system:

$$\omega_{damp}^2 = \frac{1}{2} (\Omega_{res}^2 + \omega_0^2)$$

$$\omega_{damp} = \sqrt{\omega_0^2 - \left(\frac{b}{2m}\right)^2}$$

$$\omega_{damp}^2 = \omega_0^2 - \left(\frac{b}{2m}\right)^2$$

$$x = A e^{-\frac{b}{2m}t} \cos(\omega_{damp}x + \phi)$$

- 2b) (5 points) Convince the grader that the relationship in part a holds for all simple harmonic oscillators.

- 2c) (10 points) On the way to class, you spot an abandoned bird's nest sitting in a low-hanging, more-or-less horizontal branch. You displace the end by some small amount and release it, and note that a) the tip makes about f_1 complete cycles every second, and b) it takes about N complete cycles for the amplitude of the branch's vibrations to drop to half its initial value. Find the natural frequency (f_0) for the branch/nest system. [Careful. These are f 's, not ω 's. f 's are easier to observe directly.]

$$f = f_1, \quad T = \frac{1}{f_1}, \quad \omega = 2\pi f_1$$

$$y = A e^{-\frac{b}{2m} N \cdot T} \cos \omega y$$

$$e^{-\frac{b}{2m} N \cdot \frac{1}{f_1}} = \frac{1}{2}$$

$$-\frac{b}{2m} \cdot \frac{N}{f_1} = -\ln 2$$

$$\frac{b}{2m} = \frac{\ln 2 \cdot f_1}{N}$$

$$\omega = \sqrt{\omega_0^2 - \left(\frac{b}{2m}\right)^2}$$

$$2\pi f_1 = \sqrt{\omega_0^2 - \left(\frac{\ln 2 \cdot f_1}{N}\right)^2}$$

$$4\pi^2 f_1^2 = \omega_0^2 - \left(\frac{\ln 2 \cdot f_1}{N}\right)^2$$

$$\omega_0^2 = 4\pi^2 f_1^2 + \left(\frac{\ln 2 \cdot f_1}{N}\right)^2$$

$$= \left(4\pi^2 + \frac{\ln^2 2}{N^2}\right) f_1^2$$

$$\omega_0 = f_1 \sqrt{4\pi^2 + \frac{\ln^2 2}{N^2}}$$

$$f_0 = \frac{f_1}{2\pi} \sqrt{4\pi^2 + \frac{\ln^2 2}{N^2}}$$

+10

- 2d) (5 points) Suppose we want to knock that old nest out of the tree. What would be the most effective frequency to shake the branch at?

The most effective frequency is ~~the natural frequency~~

~~$\sqrt{\omega_0^2 - \left(\frac{b}{2m}\right)^2} = 0$~~

~~$\omega_0^2 = \left(\frac{b}{2m}\right)^2$~~


~~$\omega_0 = \frac{b}{2m}$~~

~~$\frac{b}{2m} = 0$~~

~~$2m \ln 2 \cdot f_1 \cdot f_1 \sqrt{4\pi^2 + \frac{\ln^2 2}{N^2}}$~~

the fundamental frequency,

f_0



- 3a) (5 points) (Carefully) pluck a string on a violin. For a brief moment, the pluck generates noise - but that noise quickly gives way to music. Explain, in terms of physics, what is happening.

At first, the string generates noise, a mixture of all frequencies. Then, the boundary conditions filter out frequencies that are not the fundamental frequency or harmonics of that string, creating music.

+ 3

- 3b) (5 points) Derive the set of resonance frequencies for one of the open (that is, un-fingered) strings in terms of its effective length (L), the volume mass density of the material it is made of (ρ), the diameter of the string (D) and the tension in the string (T).

$$M = \frac{\rho}{4\pi D^2}$$

$$v = \sqrt{\frac{T}{M}} = \sqrt{\frac{4\pi D^2 T}{\rho}}$$

$$f_n = N \cdot \frac{v}{2L}$$

+ 3

$$= \frac{N}{2L} \sqrt{\frac{4\pi D^2 T}{\rho}}$$

- 3c) (5 points) It is not unreasonable to assume that the dominant frequency heard from an excited string will be the fundamental frequency associated with that string. One may change the fundamental frequency by pressing the string tightly into the fingerboard, effectively changing its length. Suppose you wanted to increase the fundamental frequency of a string by a factor F . Where (x) would you have to press? On a violin, the first fingered note has a frequency equal to 1.12 times the open-stringed frequency. Approximately how far up the fingerboard would you have to press to generate it?

$$Nf = \frac{N}{2 \cdot \frac{1}{N}L} \sqrt{\frac{4\pi D^2 T}{\rho}}$$

I would press at ~~at~~ $L - \frac{1}{N}L$ from top of string.

+ 3

- 3d) (10 points) Another way to change the dominant frequency you hear is to **lightly** press the string in some magic spot. Since the string is not tightly pressed, it is still able to vibrate on either side of the finger - this has the effect of imposing an intermediate node on the system, emphasizing some harmonic over the fundamental. Find each of these magic spots (x) and the frequency associated with it (in terms of the fundamental).

$$x = \frac{1}{N}L \quad f_n = N \cdot \frac{1}{2L} \sqrt{\frac{4\pi D^2 T}{\rho}}$$

+ 8

- 3e) (5 points) Bowing near the bridge can make a much nicer sound than bowing off towards the fingerboard. Explain in terms of physics why this is so.

does not reduce the length of string by too much.

+ 2