

Math 33A  
Spring 2017  
Midterm Exam 1  
4/24/2017  
Time Limit: 50 Minutes

Name (Print): Timothy Marjono  
Section: 1B

This exam contains 9 pages (including this cover page) and 4 problems. Check to see if any pages are missing. Print your name legibly where requested on the top of this page, and print your initials on the top of every page, in case the pages become separated.

You should show your work clearly and concisely. If you need more space, use the back of the pages; clearly indicate when you have done this.

Draw a box around your final answer for each problem.

You may *not* use your books, notes, or any calculator on this exam.

Do not write in the table to the right.

Problem	Points	Score
1	30	30
2	25	13
3	25	17
4	20	15
Total:	100	75

70  
00

$\frac{211}{1} \rightarrow -\frac{11}{2} \rightarrow -\frac{11}{1} \cdot \frac{2}{2}$ 
 $-11 \cdot -\frac{2}{11}$ 
717

1. Let A be the matrix  $\begin{bmatrix} 1 & -0 & 2 & 3 \\ 0 & -4 & -2 & 4 \\ 3 & 1 & 1 & -3 \\ -1 & -1 & 0 & 3 \end{bmatrix}$

(a) (20 points) Compute the reduced row echelon form (rref) of A.

$-\frac{5}{2}x = -5$

$$\begin{bmatrix} 1 & 0 & 2 & 3 \\ 0 & -4 & -2 & 4 \\ 3 & 1 & 1 & -3 \\ -1 & -1 & 0 & 3 \end{bmatrix} \begin{matrix} \\ \div (-4) \\ \\ \end{matrix}$$

$$\begin{bmatrix} 1 & 0 & 2 & 3 \\ 0 & 1 & \frac{1}{2} & -1 \\ 3 & 1 & 1 & -3 \\ -1 & -1 & 0 & 3 \end{bmatrix} \begin{matrix} \\ \\ + -3 \cdot (1) \\ + (1) \end{matrix}$$

$$\begin{bmatrix} 1 & 0 & 2 & 3 \\ 0 & 1 & \frac{1}{2} & -1 \\ 0 & 1 & -5 & -12 \\ 0 & -1 & 2 & 6 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 2 & 3 \\ 0 & 1 & \frac{1}{2} & -1 \\ 0 & 0 & -\frac{11}{2} & -11 \\ 0 & 0 & \frac{5}{2} & 5 \end{bmatrix} \begin{matrix} \\ \\ \div (-\frac{11}{2}) \\ \end{matrix}$$


$$\begin{bmatrix} 1 & 0 & 2 & 3 \\ 0 & 1 & \frac{1}{2} & -1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & \frac{5}{2} & 5 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 2 & 3 \\ 0 & 1 & \frac{1}{2} & -1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & \frac{5}{2} & 5 \end{bmatrix} \begin{matrix} + -2 \cdot (3) \\ + \frac{1}{2} \cdot (3) \\ \\ + -\frac{5}{2} \cdot (3) \end{matrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

(b) (5 points) What is the rank of  $A$ ? How do you know this?

rank  $(A) = 3$ , rank  $(A)$  is the number of leading 1's in the reduced row echelon form of  $A$ .



(c) (5 points) Find a set of vectors that spans the kernel of  $A$ .

$$\left[ \begin{array}{cccc|c} 1 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & -2 & 0 \\ 0 & 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

$$x_1 - x_4 = 0$$

$$x_2 - 2x_4 = 0$$

$$x_3 + 2x_4 = 0$$

$$x_4 = t$$

$$x_1 = x_4 = t$$

$$x_2 = 2x_4 = 2t$$

$$x_3 = -2x_4 = -2t$$

$$\begin{bmatrix} t \\ 2t \\ -2t \\ t \end{bmatrix}$$

$\rightarrow$


$$t \begin{bmatrix} 1 \\ 2 \\ -2 \\ 1 \end{bmatrix}$$

$\leftarrow$

Set of vectors that spans the kernel of  $A$

$$\begin{bmatrix} 1 \\ 2 \\ -2 \\ 1 \end{bmatrix}$$

2. Consider the mini-web shown below:



Unlike the examples you have seen on homework, this web has sites which link to themselves as well as to other sites. Also, each link has a weight between 0 and 1, such that the weights of all links leaving a site sum up to 1. The variable  $a$  is an undetermined weight with  $0 \leq a \leq 1$ .

Let  $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$  be the distribution vector of people currently viewing each website; to be a distribution vector,  $\vec{x}$  must satisfy  $x_1 + x_2 = 1$ . Assume that at each time step, everyone follows one of the links leaving their current website (or a self-link), with probabilities given by the weights on the links.

- (a) (10 points) Write down the transition matrix  $A$  such that one time step has the effect of updating  $\vec{x}$  to  $A\vec{x}$ . Give a few words of explanation. Your answer should depend on  $a$ .

$$\begin{bmatrix} \frac{1}{2} & a \\ \frac{1}{2} & 1-a \end{bmatrix}$$

10/10

for website 1,  $\frac{1}{2}$  of viewers go to website 1 and  $\frac{1}{2}$  of viewers go to website 2. This is given by first column of transition matrix, with the top number representing people who go to website 1 and bottom number representing people who go to website 2.

for website 2,  $a$  number of viewers go to website 1 and  $1-a$  viewers go to website 2. This is given by 2nd column of transition matrix, with top number representing people who go to website 1 and bottom number representing people who go to website 2.

- (b) (10 points) Find an equilibrium vector  $\vec{x}_{\text{equ}}$  for  $A$  (this means  $A\vec{x}_{\text{equ}} = \vec{x}_{\text{equ}}$ ; also,  $\vec{x}_{\text{equ}}$  must be a distribution vector, so its entries must sum to 1.). Your answer should depend on  $a$ .

$$\left[ \begin{array}{cc|c} \frac{1}{2} & a & x_1 \\ \frac{1}{2} & 1-a & x_2 \end{array} \right]$$

$$\frac{1}{2}x_1 + ax_2 = x_1$$

$$\frac{1}{2}x_1 + (1-a)x_2 = x_2$$

3/10

- (c) (5 points) For which value of  $a$  does site 2 have twice as many viewers as site 1 in the equilibrium state  $\vec{x}_{\text{equ}}$ ?

0/5

TM

3. (a) (15 points) Let  $B = \begin{bmatrix} 1 & -1 & -1 \\ 1 & 1 & -1 \\ 1 & 1 & 1 \end{bmatrix}$ . Find a matrix  $A$  with  $AB = I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ . What is  $BA$ ? You should not do another matrix multiplication to compute  $BA$ ; instead, give the answer and a conceptual justification.

$$\left[ \begin{array}{ccc|ccc} 1 & -1 & -1 & 1 & 0 & 0 \\ 1 & 1 & -1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} + -1 \cdot (1) \\ + -1 \cdot (1) \end{array}$$

$$\left[ \begin{array}{ccc|ccc} 1 & -1 & -1 & 1 & 0 & 0 \\ 0 & 2 & 0 & -1 & 1 & 0 \\ 0 & 2 & 2 & -1 & 0 & 1 \end{array} \right] \begin{array}{l} \\ \div (2) \\ \div (2) \end{array}$$

$$\left[ \begin{array}{ccc|ccc} 1 & -1 & -1 & 1 & 0 & 0 \\ 0 & 1 & 0 & -\frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 1 & 1 & -\frac{1}{2} & 0 & \frac{1}{2} \end{array} \right] + -1 \cdot (2)$$

$$\left[ \begin{array}{ccc|ccc} 1 & -1 & -1 & 1 & 0 & 0 \\ 0 & 1 & 0 & -\frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 1 & 0 & -\frac{1}{2} & \frac{1}{2} \end{array} \right] + (3)$$

$$\left[ \begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & -\frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 1 & 0 & -\frac{1}{2} & \frac{1}{2} \end{array} \right] + (2)$$

$$\left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ 0 & 1 & 0 & -\frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 1 & 0 & -\frac{1}{2} & \frac{1}{2} \end{array} \right] \leftarrow$$

$$A = \begin{bmatrix} \frac{1}{2} & 0 & \frac{1}{2} \\ -\frac{1}{2} & \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} & \frac{1}{2} \end{bmatrix}$$

$$BA = I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

A and B are square matrices, so  $AB = BA$ .

15

- (b) (10 points) Let  $C = \begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix}$ . Find a matrix  $A$  with  $AC = I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ . What is  $CA$ ? Give a geometric interpretation of the transformation  $T(\vec{x}) = CA\vec{x}$ ; be as specific as possible.

b)  $\begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$

$3 \times 2 \times (2 \times 2)$

with  $C$

$(2 \times 2)$   
matrix

+2


$T(\vec{x}) = CA\vec{x}$

4. Let  $Q$  be the linear transformation from  $\mathbb{R}^2$  to  $\mathbb{R}^2$  which is a rotation by  $\frac{\pi}{2}$  (counter-clockwise).

(a) (5 points) What are the matrices of  $Q$  and  $Q^{-1}$ ? In two separate sketches, draw the effect of  $Q$  and  $Q^{-1}$  on the unit square (the region  $0 \leq x, y \leq 1$  in the  $xy$  plane).

$$Q = \begin{bmatrix} \cos\left(\frac{\pi}{2}\right) & -\sin\left(\frac{\pi}{2}\right) \\ \sin\left(\frac{\pi}{2}\right) & \cos\left(\frac{\pi}{2}\right) \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

$$Q^{-1} = \begin{bmatrix} \cos\left(-\frac{\pi}{2}\right) & -\sin\left(-\frac{\pi}{2}\right) \\ \sin\left(-\frac{\pi}{2}\right) & \cos\left(-\frac{\pi}{2}\right) \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$



(b) (5 points) For any real number  $a$ , let  $T_a(\vec{x}) = \begin{bmatrix} 2 & 0 \\ 0 & a \end{bmatrix} \vec{x}$ . What is the matrix of  $Q \circ T_a \circ Q^{-1}$ ?

Your answer should depend on  $a$ .

$$T_a(\vec{x}) = \begin{bmatrix} 2 & 0 \\ 0 & a \end{bmatrix}$$


$$Q \circ T_a \circ Q^{-1} = \begin{bmatrix} \cos\left(\frac{\pi}{2}\right) & -\sin\left(\frac{\pi}{2}\right) \\ \sin\left(\frac{\pi}{2}\right) & \cos\left(\frac{\pi}{2}\right) \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & a \end{bmatrix} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} a & 0 \\ 0 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 0 & -a \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$



- (c) (5 points) Draw the effect of  $Q \circ T_a \circ Q^{-1}$  on the unit square in the case  $a = -1$ .



$$\begin{bmatrix} -1 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -1 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

- (d) (5 points) Let  $\begin{bmatrix} x(a) \\ y(a) \end{bmatrix} = Q \circ T_a \circ Q^{-1} \left( \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right)$ . Give explicit formulas for  $x(a)$  and  $y(a)$  as functions of  $a$ , and draw the graphs of these functions. For which value(s) of  $a$  is  $Q \circ T_a \circ Q^{-1} \left( \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right)$  a scalar multiple of  $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ?

