

# MATH 33A Midterm 2

RYAN DALY

TOTAL POINTS

**43 / 43**

QUESTION 1

## 1 Problem 1 7 / 7

✓ - 0 pts Correct

- 2 pts Incorrect orthonormal basis for the x-z plane
- 2 pts Incorrect Projection Formula
- 1 pts Incorrect Projection. It should be (1,0,3)
- 2 pts Incorrect projection vector drawing

QUESTION 2

## 2 Problem 2 10 / 10

- + 1 pts Transforms  $v_1$  correctly [no credit if T is applied, as this is the wrong vector]
- + 1 pts Transforms  $2v_1$  correctly [no credit if T is applied, as this is the wrong vector]
- + 3 pts Transforms  $e_2$  correctly [no credit if T is applied, as this is the wrong vector]
- 1 pts Computational error
- + 2 pts Transforms  $Te_2$  instead of  $e_2$  [need to specifically find the B-coordinates of  $Te_2$ , which are  $(7/5, -3/5)$  - no points for computing  $Ae_2$ ,  $A^{-1}e_2$ ,  $Be_2$ , or  $B^{-1}e_2$ ]
- ✓ + 5 pts All transforms correct
- + 3 pts Used S instead of  $S^{-1}$  for the transforms
- ✓ + 5 pts Finds B-matrix correctly
- 1 pts Computational error
- + 2 pts Gives formula for B
- + 0 pts No credit

QUESTION 3

## 3 Problem 3 6 / 6

✓ - 0 pts Correct

- 2 pts Incorrect solutions to the matrix. That is, finding the kernel.
- 2 pts Incorrect basis for the kernel.
- 2 pts Incorrect basis for the image.

QUESTION 4

## 4 Problem 4 10 / 10

✓ + 10 pts Correct

- + 5 pts Rubric Criterion 1: Correctly Calculates Q
- + 5 pts Rubric Criterion 2: Correctly Calculates R
- + 3 pts Potential Partial Credit: Student makes a small mistake in the calculation of R or Q, i.e. gets one entry incorrect.
- + 0 pts Incorrect/ insufficient progress

QUESTION 5

## Problem 5 10 pts

### 5.1 2 / 2

- ✓ + 2 pts Correct, since the columns vectors span a subspace of  $\mathbb{R}^4$ .
- + 0 pts Wrong, if they were, they would span a 5-dimensional subspace of a 4-dimensional space!
- + 0 pts no answer

### 5.2 2 / 2

- ✓ + 2 pts Correct, as  $5 = \dim(V) + \dim(V_{\text{orth}})$
- + 0 pts Wrong. For that to be true you would need to have  $\mathbb{R}^n$  with n an even number instead of  $\mathbb{R}^5$

### 5.3 2 / 2

- ✓ + 2 pts Correct, since we can take S to be the identity matrix, and then  $AS=SA$
- + 0 pts Wrong
- + 0 pts no answer

### 5.4 2 / 2

- ✓ + 2 pts Correct, since by the rank-nullity theorem the dimension of the kernel and image sum up to 5. The image can be at most 3-dimensional, thus the kernel has to be at least 2-dimensional.

+ 0 pts Wrong. For this to be true the image would need to be a 4-dimensional subspace of a 3-dimensional space.

+ 0 pts no answer

5.5 2 / 2

✓ + 2 pts Correct, since the kernel of the orthogonal projection is the orthogonal complement of  $V$ .

+ 0 pts Wrong

+ 0 pts no answer

## Midterm 2 (Math 33A, Fall 2019)

Your Name: Ryan DalyUCLA id: 505-416-119Date: 11/18/19

**The rules:** You can answer using a pencil or ink pen. You are allowed to use only this paper, pencil or pen, and the scratch paper provided. You should not hand the scratch paper in. No calculators. No books, no notebooks, no notes, no mobile phones, no web access. You must write your name and UCLA id. You have exactly 50 minutes.

**Points:**

Problem 1: \_\_\_\_/7

Problem 2: \_\_\_\_/10

Problem 3: \_\_\_\_/6

Problem 4: \_\_\_\_/10

Problem 5: \_\_\_\_/10

Total: \_\_\_\_\_ (out of 43)

**Problem 1** (7 points)

Let  $V$  be the  $x$ - $z$  plane in  $\mathbb{R}^3$ .

What is the orthogonal projection of  $v = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$  onto  $V$ ?

You should (i) compute the projection  $\text{proj}_V(v)$  and (ii) make a drawing of  $\text{proj}_V(v)$ .

**Solution:** orthogonal basis of  $V$  is  $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$

$$\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \quad \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 3 \end{pmatrix}$$
$$\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ 3 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 3 \end{pmatrix}$$
$$\text{proj}_V(v) = \begin{pmatrix} 1 \\ 0 \\ 3 \end{pmatrix}$$

**Solution:**

**Problem 2** (10 points)

Let  $\mathcal{B}$  be the basis of  $\mathbb{R}^2$  given by the vectors

$$v_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

and

$$v_2 = \begin{bmatrix} -1 \\ 3 \end{bmatrix}.$$

Find the  $\mathcal{B}$ -coordinates of the vectors  $v_1$ ,  $2v_1$  and  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ . Let  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be the linear transformation given by the matrix

$$A = \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix}.$$

Find the  $\mathcal{B}$ -matrix of  $T$ .

**Solution:**

$$[v_1]_{\mathcal{B}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad [2v_1]_{\mathcal{B}} = \begin{pmatrix} 2 \\ 0 \end{pmatrix} \quad [e_2]_{\mathcal{B}} = \begin{bmatrix} 1/5 \\ 1/5 \end{bmatrix}$$

$$\left[ \begin{array}{cc|c} 1 & -1 & 0 \\ 2 & 3 & 1 \end{array} \right] \xrightarrow{-2R_1} \left[ \begin{array}{cc|c} 1 & -1 & 0 \\ 0 & 5 & 1 \end{array} \right] \xrightarrow{:5} \left[ \begin{array}{cc|c} 1 & -1 & 0 \\ 0 & 1 & 1/5 \end{array} \right] \xrightarrow{+R_2} \left[ \begin{array}{cc|c} 1 & 0 & 1/5 \\ 0 & 1 & 1/5 \end{array} \right]$$

$\mathcal{B}$ -matrix of  $T$

$$B = \left[ [Tv_1]_{\mathcal{B}} \quad [Tv_2]_{\mathcal{B}} \right]$$

$$\begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 5 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} -1 \\ 3 \end{bmatrix} = \begin{bmatrix} 5 \\ 5 \end{bmatrix}$$
$$T(v_2) = \begin{bmatrix} 5 \\ 5 \end{bmatrix}$$

Solution:

$$B = \left[ \begin{bmatrix} 5 \\ 0 \end{bmatrix}_B \quad \begin{bmatrix} 5 \\ 5 \end{bmatrix}_B \right] \quad \begin{pmatrix} 1 \\ 2 \end{pmatrix} \quad \begin{pmatrix} -1 \\ 3 \end{pmatrix}$$

$$\left[ \begin{array}{cc|c} 1 & -1 & 5 \\ 2 & 3 & 0 \end{array} \right] \xrightarrow{-2I} \left[ \begin{array}{cc|c} 1 & -1 & 5 \\ 0 & 5 & -10 \end{array} \right] \div 5 \quad \left[ \begin{array}{cc|c} 1 & -1 & 5 \\ 0 & 1 & -2 \end{array} \right] \xrightarrow{+II}$$

$$\left[ \begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & -2 \end{array} \right] \quad B = \begin{bmatrix} 3 \\ -2 \end{bmatrix} \begin{bmatrix} 5 \\ 5 \end{bmatrix}_B$$

$$\left[ \begin{array}{cc|c} 1 & -1 & 5 \\ 2 & 3 & 5 \end{array} \right] \xrightarrow{-2I} \left[ \begin{array}{cc|c} 1 & -1 & 5 \\ 0 & 5 & -5 \end{array} \right] \div 5 \quad \left[ \begin{array}{cc|c} 1 & -1 & 5 \\ 0 & 1 & -1 \end{array} \right] \xrightarrow{+II}$$

$$\left[ \begin{array}{cc|c} 1 & 0 & 4 \\ 0 & 1 & -1 \end{array} \right] \quad B = \begin{bmatrix} 3 & 4 \\ -2 & -1 \end{bmatrix}$$

**Problem 3** (6 points)

Let

$$A = \begin{bmatrix} 1 & 0 & -1 \\ 2 & 0 & -2 \\ -1 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

Find a basis for the image of  $A$  and a basis for the kernel of  $A$ .

**Solution:**

$$\begin{bmatrix} 1 & 0 & -1 \\ 2 & 0 & -2 \\ -1 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{array}{l} -2I \\ +I \\ \end{array} \begin{bmatrix} 1 & 0 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

column 2 and 3 are redundant

$\text{im}(A) = \text{span}\left(\begin{pmatrix} 1 \\ 2 \\ -1 \\ 0 \end{pmatrix}\right)$  so a basis of  $\text{im}(A)$   
is  $\left(\begin{pmatrix} 1 \\ 2 \\ -1 \\ 0 \end{pmatrix}\right)$

$$\left[ \begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

for  $x_3 = r, x_4 = t$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} r \\ t \\ r \end{bmatrix} = r \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + t \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

a basis of  $\text{ker}(A)$  is  $\left(\begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}\right) \left(\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}\right)$

$$\dim(\text{ker}(A)) + \dim(\text{im}(A)) = 1 + 2 = 3$$



**Solution:**

**Problem 4** (10 points)

Compute the QR decomposition of the following matrix.

$$M = \begin{bmatrix} -1 & -1 \\ 1 & 3 \\ -1 & -1 \\ 1 & 3 \end{bmatrix}$$

**Solution:**

$$\begin{bmatrix} -1 & -1 \\ 1 & 3 \\ -1 & -1 \\ 1 & 3 \end{bmatrix} \quad R_{11} = \sqrt{4} = 2 \quad u_1 = \frac{1}{2} \begin{bmatrix} -1 \\ -1 \\ 1 \\ 1 \end{bmatrix}$$
$$R_{12} = u_1 \cdot v_2 = \frac{1}{2} \begin{pmatrix} -1 \\ -1 \\ 1 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 3 \\ -1 \\ 3 \end{pmatrix} = \frac{1}{2} (1 + 3 + 1 + 3) = \frac{1}{2} 8 = 4$$
$$v_2^\perp = \begin{pmatrix} -1 \\ 3 \\ -1 \\ 3 \end{pmatrix} - 4 \left( \frac{1}{2} \right) \begin{pmatrix} -1 \\ -1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} -1 \\ 3 \\ -1 \\ 3 \end{pmatrix} - \begin{pmatrix} -2 \\ 2 \\ 2 \\ 2 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ -3 \\ 1 \end{pmatrix}$$
$$R_{22} = \sqrt{4} = 2 \quad u_2 = \frac{1}{2} \begin{pmatrix} 1 \\ 1 \\ -3 \\ 1 \end{pmatrix}$$

$$QR = \frac{1}{2} \begin{bmatrix} -1 & 1 \\ -1 & 1 \\ 1 & -3 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 4 \\ 0 & 2 \end{bmatrix}$$

$$v_2 = \frac{1}{2} \begin{pmatrix} -4+2 \\ 4+2 \\ -4+2 \\ 4+2 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} -2 \\ 6 \\ -2 \\ 6 \end{pmatrix} = \begin{pmatrix} -1 \\ 3 \\ -1 \\ 3 \end{pmatrix}$$

$$v_4 = 2 \left( \frac{1}{2} \right) \begin{pmatrix} -1 \\ -1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} -1 \\ -1 \\ 1 \\ 1 \end{pmatrix}$$

**Solution:**

**Problem 5** (10 points total; 2 points each)

Answer the following questions with true or false.

max rank = 4

1. Let  $A$  be an arbitrary  $4 \times 5$  matrix (i.e., with 4 rows and 5 columns). The column vectors of  $A$  cannot be linearly independent.

True

2. There exists a subspace  $V$  of  $\mathbb{R}^5$  so that  $V$  and its orthogonal complement  $V^\perp$  have the same dimension.

False

3. Any square matrix  $A$  is similar to itself.

True

4. There exists a linear transformation  $T: \mathbb{R}^5 \rightarrow \mathbb{R}^3$  with kernel of dimension 1.

False

5. Let  $V$  be a subspace of  $\mathbb{R}^n$ , and  $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$  the orthogonal projection onto  $V$ . Let  $v$  be a vector in the kernel of  $T$ . Then  $v \cdot w = 0$  for all vectors  $w$  in  $V$ .

True

3  $\left[ \begin{array}{c} \overbrace{\hspace{2cm}}^5 \\ \hspace{2cm} \end{array} \right]$   
max rank = 3  
min ker = 2

$V^\perp$  perpendicular to  $V$

Use this sheet of paper if you need additional space.

Use this sheet of paper if you need additional space.