


Full Name \_\_\_\_\_ UID \_\_\_\_\_

Yanli Liu	1A T 9:00A-9:50A MS 6229 1B R 9:00A-9:50A Boelter 5419
Nicholas Boschert	1C T 9:00A-9:50A Boelter 5280 1D R 9:00A-9:50A Boelter 5280
Gyu Eun Lee	1E T 9:00A-9:50A Boelter 5272 1F R 9:00A-9:50A Boelter 5272

Section	1	F
---------	---	---

- Do not open this exam packet until you are told that you may begin.
- Turn off all electronic devices and and put away all items except for a pen/pencil and an eraser.
- Remove hats and sunglasses.
- No phones, calculators, smart-watches or electronic devices of any kind allowed for any reason, including checking the time.
- If you have a question, raise your hand and one of the proctors will come to you. We will not answer any mathematical questions except possibly to clarify the wording of a problem.
- Quit working and close this packet when you are told to stop.



Page:	1	2	3	4	5	6	7	8	Total
Points:	6	4	8	7	6	8	5	6	50
Score:	6	4	4	7	6	8	5	6	46

Part I: Multiple choice. Please write your answers (A, B, C, ...) in the boxes on the right.

1. (3 points) Determine values of the constants  $a$  and  $b$  that make the differential equation exact.

$$(3x^2y - bx^5y^3) dx + (ax^3 + x^6y^2) dy = 0 \quad (1)$$

- A.  $a = 2, b = -1$
- B.  $a = 1, b = -2$
- C.  $a = 2, b = 1$
- D.  $a = -1, b = 2$
- E. None of the above.

$$\frac{\partial P}{\partial y} = 3x^2 - 3bx^5y^2$$

$$\frac{\partial Q}{\partial x} = 3ax^2 + 6x^5y^2$$

$$a = 1, b = -2$$

B

2. (3 points) Consider the initial value problem


$$y' = (y^2 - 1) \sin^2(ty), \quad y(0) = 2.$$

Which of the following is true?

- A.  $-1 < y(t) < 1$  for all  $t$  for which  $y$  is defined.
- B.  $y(t) < \sin^2(t)$  for all  $t$  for which  $y$  is defined.
- C.  $y(t) > 1$  for all  $t$  for which  $y$  is defined.
- D.  $y(t) > t^2 - 1$  for all  $t$  for which  $y$  is defined.
- E. None of the above.


$y = 1$

$$\text{LHS} = 0 \quad \text{RHS} = (0) \sin^2(y) = 0$$



C

3. (4 points) The slope field below corresponds to which differential equation?



- A.  $y' = \frac{t^2}{1-y^2}$
- B.  $y' = t^2$
- C.  $y' = 4y$
- D.  $y' = ty$
- E.  $y' = \frac{y^2}{2t+1}$

A

4. (4 points) Determine if the equation

$$\left(\frac{y}{x} + 8x\right) dx + (\ln x - 3) dy = 0 \quad (2)$$

is exact (you may assume that we are working in a rectangle  $R$  in the plane such that  $x > 0$  for all  $(x, y)$  in  $R$ ). If it is exact, find the solution.

A.  $F(x, y) = y \ln x + 4x^2 - 3y$

$$\frac{\partial P}{\partial y} = \frac{1}{x}$$

B.  $F(x, y) = -\frac{y}{x^2} + \frac{1}{x} + 8$

C.  $-\frac{y}{x^2} + \frac{1}{x} + 8 = C$

$$\frac{\partial Q}{\partial x} = \frac{1}{x}$$

D.  $y \ln x + 4x^2 - 3y = C$

E. The equation is not exact.

$$\frac{\partial F}{\partial x} = \frac{y}{x} + 8x$$

$$F(x, y) = y \ln x - 3y + 4x^2 = C$$

$$\frac{\partial F}{\partial y} = \ln x - 3 \Rightarrow F = \int (\ln x - 3) dy$$

$$= y \ln x - 3y + g(x)$$

$$\frac{\partial F}{\partial x} = \frac{y}{x} + g'(x) = \frac{y}{x} + 8x$$

$$g'(x) = 8x$$

$$g(x) = 4x^2$$

D

5. (4 points) Which of the following integrating factors is suitable for the differential equation

$$(x + 2) \sin y dx + x \cos y dy = 0? \quad (3)$$

A.  $e^{\cos x}$

B.  $\sin x$

C.  $x e^x$

D.  $1 + \frac{1}{x}$

E. None of the above.

$$x \sin y + 2 \sin y$$

$$x \cos y + 2 \cos y \quad \cos y$$

$$\frac{1}{x \cos y} (x \cos y + \cos y)$$

$$\frac{(x+1)(\cos y)}{x(\cos y)} = \frac{x+1}{x} = 1 + \frac{1}{x}$$

D

6. (4 points) The function  $\mu(x, y) = \frac{1}{x^2 + y^2}$  is an integrating factor for the equation

$$(x^2 + y^2 - x)dx - y dy = 0. \quad (4)$$

Use this to solve the differential equation. You may assume that we are working in a rectangle  $R$  which does not contain the point  $(0, 0)$ .

- A.  $F(x, y) = x - \arctan(x^2 + y^2)$
- B.  $F(x, y) = x - \frac{1}{2} \ln(x^2 + y^2)$
- C.  $x - \frac{1}{2} \ln(x^2 + y^2) = C$
- D.  $x - \arctan(x^2 + y^2) = C$
- E. None of the above

$$\left(1 - \frac{x}{x^2 + y^2}\right) dx - \frac{y}{x^2 + y^2} dy = 0$$

$$\frac{\partial F}{\partial x} = 1 - \frac{x}{x^2 + y^2}$$

$$\frac{\partial F}{\partial y} = -\frac{y}{x^2 + y^2} \Rightarrow F = \int \frac{-y}{x^2 + y^2} dy$$

$u = x^2 + y^2$   
 $du = 2y dy$

$$= -\frac{1}{2} \int \frac{du}{u}$$

$$= -\frac{1}{2} \ln(x^2 + y^2) + g(x)$$

C

7. (3 points) True or false: there exists a differential equation of the form  $y' = f(t, y)$  such that  $f$  has continuous partial derivatives on a rectangle  $R$  containing  $(0, 0)$  and such that

$$y_1 = 2t \quad \text{and} \quad y_2 = 3t \quad (5)$$

are both solutions in  $R$ .

- A. False. The existence theorem forbids it.
- B. False. The uniqueness theorem forbids it.
- C. True. The existence theorem guarantees it.
- D. True. The uniqueness theorem guarantees it.

B

Part II: Free Response. Write up a full solution for each problem. A correct answer with an incomplete or incorrect solution will not receive full credit.

8. (6 points) Find the general solution of the linear equation

$$y' - 2y = 4t^3 e^{2t}.$$

Box your answer

$$y' = 2y + 4t^3 e^{2t}$$

$$\frac{dy_h}{dt} = 2y_h$$

$$\int \frac{dy_h}{y_h} = \int 2 dt$$

$$\ln|y_h| = 2t + C_1$$

$$y_h = Ae^{2t} = v(t)e^{2t} \quad (A = \pm e^{C_1})$$

$$y' = v'(t)e^{2t} + \cancel{v(t)2e^{2t}} = \cancel{2(v(t)e^{2t})} + 4t^3 e^{2t}$$

$$v(t) = \int 4t^3 dt$$


$$= t^4 + C_2$$

$$y(t) = e^{2t} t^4 + C_2 e^{2t}$$

Gen. soln.:

$$y(t) = t^4 e^{2t} + C e^{2t}$$

9. (8 points) Consider two tanks, labeled tank A and tank B. Tank A contains 150 gal of solution in which is dissolved 12 lbs of salt. Tank B contains 300 gal of solution in which is dissolved 28 lbs of salt. Pure water flows into tank A at a rate of 4 gal/s. There is a drain at the bottom of tank A. The solution leaves tank A via this drain at a rate of 4 gal/s and flows immediately into tank B at the same rate. A drain at the bottom of tank B allows the solution to leave tank B at a rate of 1 gal/s. **Set up, but do not evaluate**, a system of differential equations involving the variables  $x$  (amount of salt in tank A),  $y$  (amount of salt in tank B), and  $t$  (time in seconds). Do not forget to include any initial conditions. Box your entire answer



$$A: \frac{dx}{dt} = \left( 0 \frac{\text{lb}}{\text{gal}} \cdot 4 \frac{\text{gal}}{\text{s}} \right) - \left( \frac{x \text{ lb}}{150 \text{ gal}} \cdot 4 \frac{\text{gal}}{\text{s}} \right)$$

$$= 0 - \frac{4x \text{ lb}}{150 \text{ s}}$$

$$B: \frac{dy}{dt} = \left( \frac{4x \text{ lb}}{150 \text{ s}} \right) - \left( \frac{y \text{ lb}}{300+3t \text{ gal}} \cdot 1 \frac{\text{gal}}{\text{s}} \right)$$

$$= \frac{4x}{150} - \frac{y}{300+3t}$$

$$\frac{dx}{dt} = - \frac{4x}{150}$$

$$\frac{dy}{dt} = \frac{4x}{150} - \frac{y}{300+3t}$$

Initial conditions:

$$x(0) = 12$$

$$y(0) = 28$$

$V_A = 150 \text{ gal}$

$$V_B = V_0 + (4 \text{ gal/s} - 1 \text{ gal/s})t$$

$$= 300 \text{ gal} + 3 \text{ gal/s} t$$

Rate = rate in - rate out

10. (5 points) The differential equation

$$y'' + 2t(y')^2 = 0$$

is an example of a second order differential equation. The change of variable  $v = y'$  (so that  $v' = y''$ ) turns this equation into a first-order separable equation. Using this change of variable, find the particular solution which satisfies

$$y(1) = \frac{\pi}{2}, \quad y'(0) = 1.$$

Box your answer

$$y'' + 2t(y')^2 = 0$$

$$v' + 2tv^2 = 0$$

$$v' = -2tv^2$$

$$\int \frac{dv}{v^2} = \int -2t \, dt$$

$$-\frac{1}{v} = -t^2 + C_1$$

$$v = \frac{1}{t^2 - C_1}$$

$$y' = \frac{1}{t^2 - C_1}$$

$$y'(0) = 1 = \frac{1}{0 - C_1}$$
$$1 = \frac{1}{-C_1}$$
$$C_1 = -1$$

$$y' = \frac{1}{t^2 + 1}$$

$$y = \int \frac{1}{t^2 + 1} \, dt$$

$$y = \arctan t + C_2$$

$$y(1) = \frac{\pi}{2} = \arctan(1) + C_2$$

$$\frac{\pi}{2} = \frac{\pi}{4} + C_2$$

$$C_2 = \frac{\pi}{4}$$

$$y(t) = \arctan t + \frac{\pi}{4}$$



11. (6 points) The differential equation

$$y^2 y' = -y^3 + 3e^{-t} \quad (6)$$

is an example of a Bernoulli equation with degree  $n = -2$ .

- (a) The change of variables  $z = y^3$  turns this equation into a first-order linear differential equation. Using this, write the linear equation above in the form  $z' = a(t)z + f(t)$ . (I will not be grading your work for this problem, just your answer). Box your answer.

$$z = y^3$$

$$\frac{dz}{dy} = \frac{d(y^3)}{dy}$$

$$\frac{dz}{dy} = 3y^2$$

$$\frac{dy}{dt} = y' = -\frac{y^3}{y^2} + \frac{3e^{-t}}{y^2} = -y + \frac{3e^{-t}}{y^2}$$

$$\frac{dz}{dy} \cdot \frac{dy}{dt} = 3y^2 \left( -y + \frac{3e^{-t}}{y^2} \right)$$

$$\frac{dz}{dt} = -3y^3 + 9e^{-t} = -3z + 9e^{-t}$$

$$z' = -3z + 9e^{-t}$$

- (b) Solve the differential equation. Your answer should be in the form  $y(t) = \dots$

$$\frac{dz_h}{dt} = -3z_h$$

$$\int \frac{dz_h}{z_h} = \int -3 dt$$

$$\ln|z_h| = -3t + C \quad (A = \pm e^C)$$

$$z_h = A e^{-3t} = v(t) e^{-3t}$$

$$z' = v'(t) e^{-3t} + v(t) (-3) e^{-3t} = -3(v(t) e^{-3t}) + 9e^{-t}$$

$$v'(t) = \int 9e^{2t} dt$$

$$v(t) = \frac{9}{2} e^{2t} + C_2$$

$$z(t) = \frac{9}{2} e^{-t} + C_2 e^{-3t}$$

$$y^3 = \frac{9}{2} e^{-t} + C_2 e^{-3t}$$

$$y(t) = \sqrt[3]{\frac{9}{2} e^{-t} + C_2 e^{-3t}}$$