

33B Midterm 2

TOTAL POINTS

38 / 40

QUESTION 1

auto. Equation 11 pts

1.1 Phase Line 3 / 3

- ✓ - 0 pts Correct
- 1 pts arrows missing/wrong
- 1 pts graph wrong somewhere
- 1 pts missing scale
- 3 pts not the phase line
- 1 pts too many zero's in the graph

1.2 Eq. Solutions 3 / 3

- ✓ - 0 pts Correct
- 1 pts -3 no conclusion
- 1 pts 2 is stable
- 1 pts 5 is unstable
- 2 pts Stable/unstable?

1.3 Graph sketch 2 / 2

- ✓ - 0 pts Correct
- 2 pts no solution/wrong solution
- 1 pts graph between -3 and 2 wrong/missing
- 1 pts graph between 2 and 5 wrong/missing

1.4 particular solution 3 / 3

- ✓ + 3 pts Correct
- + 1 pts No
- + 1 pts Uniqueness theorem can be applied
- + 0 pts wrong/no answer
- + 1 pts cannot cross the equilibrium solution $y(t) = 2$

QUESTION 2

Existence and Uniqueness 8 pts

2.1 Apply? Rectangle? 5 / 5

- ✓ + 2 pts continuous

✓ + 2 pts derivative continuous

✓ + 1 pts rectangle

+ 0 pts no points

2.2 $x_0(2)=5?$ 3 / 3

✓ + 1 pts Correct

✓ + 2 pts justification

+ 0 pts no points

QUESTION 3

3 Particular Solution 6 / 6

✓ - 0 pts Correct

- 1 pts Mixed up a minus sign

- 3 pts Didn't try the right guess (ae^{3t})

- 6 pts Didn't attempt method of undetermined coefficients.

- 1 pts Incorrect arithmetic in finding constant.

- 1 pts Incorrect multiplication

- 1 pts Put constant in solution

- 3 pts Forgot to include an undetermined coefficients in MOC.

QUESTION 4

2. order equation constant coefficients 5 pts

4.1 verify solutions 3 / 3

✓ - 0 pts Correct

- 2 pts Didn't explicitly check boundary conditions

- 1 pts Only checked one boundary condition

- 1 pts Didn't correctly check that they satisfy the ODE.

4.2 existence and uniqueness? 0 / 2

- 0 pts Correct

- 2 pts Didn't understand that solution was non-

unique.

✓ - **2 pts** Didn't state that the boundary conditions being defined at different times renders the existence and uniqueness theorem irrelevant.

- **1 pts** Not clear if you actually meant that the "initial" conditions are defined at different times.

QUESTION 5

2. order equation 7 pts

5.1 verify solutions 4 / 4

✓ - **0 pts** Correct

- **2 pts** incorrect calculation

- **4 pts** incorrect calculation

5.2 fundamental set 3 / 3

✓ - **0 pts** Correct

- **1 pts** conclusion is incorrect,

- **1 pts** some work, calculation incorrect,

- **3 pts** conclusion incorrect, wrong calculation

- **2 pts** some work

QUESTION 6

6 planar system 3 / 3

✓ - **0 pts** Correct

- **2 pts** incorrect, but some work

- **1 pts** minor mistake

- **3 pts** no work

79

MIDTERM 2

11/16/2018

Math33B
Nadja Hempel
nadja@math.ucla.edu

Name: [REDACTED]

UID: [REDACTED]

section: [REDACTED]

Problem	Points	Score
1	11	
2	8	
3	6	
4	5	
5	7	
6	3	
Total	40	

Instructions


- (1) This exam has 6 problems. Make sure you have all pages.
- (2) Enter your name, SID number, and discussion section on the top of this page.
- (3) Use a **PEN** to record your final answers.
- (4) If you need **more space**, use the extra page at the end of the exam.
- (5) **NO** Calculators, computers, books or notes of any kind are allowed.
- (6) Show your work. Unsupported answers will not receive full credit.
- (7) Good Luck!

Exercise 1. (11pt)

Consider the autonomous first-order differential equation.

$$y' = (y + 3)^2(y - 2)(y - 5)$$


- (1) Draw the phase line. (3pt)



- (2) What are the equilibrium solutions? Which are stable, and which are unstable?
-
- (3pt)

 -3 : undetermined 2 : stable 5 : unstable

- (3) Sketch the graph of at least one solution between each pair of adjacent equilibrium solutions. (2pt)



- (4) Let $y_p(t)$ be a particular solution to the equation which satisfies $y_p(0) = 0$. Is it possible that $y_p(2) = 2$? Justify your answer. (3pt)


No: $y(t) = 2$ is an equilibrium solution and so is $y(t) = -3$. According to the uniqueness theorem, only one solution can exist with $y_p(2) = 2$, which is $y(t) = 2$, so no others can cross it. The solution corresponding to $y_p(0) = 0$ is some solution between $y(t) = -3$ and $y(t) = 2$.

Exercise 2. (8pt) Consider the differential equation

$$\frac{dx}{dt} = \frac{\sqrt{x^2 - 4}}{t^2}$$

- (1) Can you apply the existence and uniqueness theorem to the initial value problem $x_0(1) = 6$? Justify your answer and give the biggest rectangle in which you can apply it containing the given point (if it exists). (5 pt)

$$f(x, t) = \frac{\sqrt{x^2 - 4}}{t^2} \quad \text{cont. } x \geq 2 \quad x \leq -2 \\ t \neq 0$$



$$\frac{\partial f}{\partial x} = \frac{2x}{2t^2\sqrt{x^2 - 4}} = \frac{x}{t^2\sqrt{x^2 - 4}} \quad \text{cont. } x > 2 \quad x < -2 \\ t \neq 0$$

The existence and uniqueness thm. applies to the largest rect. on which $f(x, t)$ and $\frac{\partial f}{\partial x}$ are continuous. In this case, for $x_0(1) = 6$,

$$R : \quad x \in (2, \infty) \\ t \in (0, \infty)$$

- (2) Can $x_0(2) = 5$ ($x_0(t)$ is the solution to the initial value problem in part 1))?(1pt)
Justify your answer. (2pt)

No, $\frac{dx}{dt}$ is always ≥ 0 , so
an equation $x_0(t)$ with a point
 $x_0(1) = 6$ will have $x_0(2) \geq 6$.

Exercise 3. (6pt) Find a particular solution to the following differential equation

$$3y'' + 2y' - y = -4e^{3t}.$$

Try: $y = Ce^{3t}$
 $y' = 3Ce^{3t}$
 $y'' = 9Ce^{3t}$

$$27Ce^{3t} + 6Ce^{3t} - Ce^{3t} = -4e^{3t}$$

$$32Ce^{3t} = -4e^{3t}$$

$$\Rightarrow C = -\frac{1}{8}$$

$$y_p(t) = -\frac{1}{8} e^{3t}$$

Exercise 4. (5pt) Consider the following problem:

$$y'' + y = 0 \quad y(0) = 0 \quad y'(\pi/2) = 0$$

(1) Show that $y(t) = C \cdot \sin(t)$ is a solution for any constant C . (3pt)

$$y'(t) = C \cos t$$

$$y''(t) = -C \sin t$$


$$y'' + y = -C \sin t + C \sin t = 0 \quad \checkmark$$

$$y(0) = C \sin(0) = 0 \quad \checkmark$$

$$y'(\pi/2) = C \cos(\pi/2) = 0 \quad \checkmark$$

(2) Why does this not violate the 2. order existence and uniqueness theorem? (2pt)

This is a homogeneous eqn, discontinuous at $y=0$, so the existence & uniqueness theorem only applies on intervals where $y \neq 0$.



Exercise 5. (8pt) Consider the differential equation

$$y'' + \frac{1+x}{x}y' - \frac{1}{x}y = 0$$

- (1) Check that $1+x$ and $\frac{2x^2+6x+4}{x-2}$ are solutions to the above equation. (4pt)

$$y_1 = 1+x$$

$$y_1' = 1$$

$$y_1'' = 0$$

$$0 + \frac{1+x}{x}(1) - \frac{1}{x}(1+x) = 0 \quad \checkmark$$

$$y_2 = \frac{2x^2+6x+4}{x+2} = \frac{(2x+2)(x+2)}{x+2} = 2x+2$$

Removable discant?
 $x \neq -2$

$$y_2' = 2$$

$$y_2'' = 0$$

$$0 + \frac{1+x}{x}(2) - \frac{1}{x}(2x+2) = 0 \quad \checkmark$$

(2) Do they form a fundamental set of solutions? (1pt) Justify your answer. (2pt)

No, they are not linearly independent.

$$\frac{y_2}{y_1} = \frac{2x+2}{x+1} = 2$$

$$\Rightarrow y_2 = 2y_1$$

↑
constant

Exercise 6. (3pt)

Consider the second order equation

$$y'' - 2e^t y' - \tan(t)y = \sqrt{t^2 + 1}.$$

Write this equations as a planar system of first-order equations.

$$y'' = 2e^t y' + \tan(t)y + \sqrt{t^2 + 1}.$$

$$x = y'$$

$$x' = y''$$

$$\begin{array}{l} x' = 2e^t x + \tan(t)y + \sqrt{t^2 + 1} \\ x = y' \end{array}$$

Extra page

Extra page