

Midterm 1

Last Name: _____

First Name: _____

Student ID: _____

Section:

Tuesday:

Thursday:

1A

1B

TA: Albert Zheng

1C

1D

TA: Benjamin Spitz

1E

1F

TA: Eilon Reisin-Tzur

Instructions: Do not open this exam until instructed to do so. You will have 50 minutes to complete the exam. Please print your name and student ID number above, and circle the number of your discussion section. **You may not use calculators**, books, notes, or any other material to help you. Please make sure **your phone is silenced** and stowed where you cannot see it. Remember that you are bound by a conduct code, and that you may not look at anyone's paper or let anyone look at your paper, nor may you work on the exam after it has ended. There is some scratch paper at the back of the exam, if you do work there indicate so on the relevant question. Do not write on the backs of the pages. Please circle or box your final answers.

Please get out your id and be ready to show it when you turn in your exam.

Please do not write below this line.

Question	Points	Score
1	10	
2	10	
3	10	
4	10	
5	10	
Total:	50	

1. (10 points) Circle the correct answer (only one answer is correct for each question)


1. The relation on the integers defined by xRy if $x - y$ is odd is:

- (a) an equivalence relation
- (b) symmetric, but not transitive or reflexive
- (c) reflexive, but not symmetric or transitive
- (d) symmetric and transitive, but not reflexive

2. If $f : X \rightarrow Y$ is an one-to-one function and $g : Y \rightarrow Z$ is an onto function, then $g \circ f$ is necessarily:

- (a) one-to-one
- (b) onto
- (c) a bijection
- (d) not necessarily one-to-one or onto

For every x there is a different y
and for every z there is some y



3. If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are functions and $g \circ f$ is a bijection, then:

- (a) f must be one-to-one and g must be onto
- (b) f and g must both be bijections
- (c) f must be onto
- (d) g must be one-to-one

$g(f(x))$

4. The function $f : \{a, b\}^* \times \{a, b\}^* \rightarrow \{a, b\}^*$ (X^* is the set of strings in X) defined by $f((\alpha, \beta)) = \alpha\beta$ is:

- (a) onto and one-to-one
- (b) one-to-one but not onto
- (c) neither one-to-one nor onto
- (d) onto but not one-to-one

5. For X a set with n elements, how many relations are there on X that are both partial orders and equivalence relations?

- (a) more than 1 and less than n
- (b) more than n
- (c) exactly n
- (d) exactly 1

Need to be both
Symmetric & antisymmetric
reflexive
transitive



2. In this question write down your answer, no need for any justification.

- (a) (2 points) Give an example of a functions $f : X \rightarrow Y$, $g : Y \rightarrow Z$ so that g is onto but $g \circ f$ is not onto.

let X, Y, Z
be \mathbb{R}

$$f(x) = x^2 \quad g(y) = y \quad \leftarrow \text{onto}$$

$$g(f(x)) = x^2 \quad \leftarrow \text{not onto, only positive outputs!}$$

- (b) (2 points) What is $\sum_{i=0}^{100} i$? Feel free to write your answer as the product or sum of a few numbers.

$$0+1+2+3+4+5$$

$$0 \quad 1 \quad 3 \quad 6 \quad 10 \quad 15$$

$$n \cdot \left(\frac{n+1}{2}\right) = 100 \cdot \left(\frac{101}{2}\right) = 50 \cdot 101$$

- (c) (2 points) List elements of the equivalence relation on $\{a, b, c, d, e\}$ determined by the partition $\{\{a, b\}, \{c\}, \{d, e\}\}$.

$$[a] = \{a, b\} \quad [c] = \{c\} \quad [d] = \{d, e\}$$


$$[b] = \{a, b\} \quad [e] = \{d, e\}$$

- (d) (2 points) If X is a set with n elements where $n \geq 1$, how many onto functions are there from X to $\{0, 1\}$?

$$n(n-1)$$

- (e) (2 points) Give an example of a relation on $X = \{a, b, c\}$ that is reflexive and symmetric but not transitive.

$$\{(a, a), (b, b), (c, c), (a, b), (b, a), (b, c), (c, b)\}$$



3. Let $X = \mathbb{Z} \times \mathbb{Z}_{>0}$ (i.e. ordered pairs of integers where the second integer is positive), and define a relation Q on X by $(a, b)Q(c, d)$ if $ad = bc$.

(a) (8 points) Show that Q is an equivalence relation on X .

Reflexive

$$(a, b)Q(a, b) \quad ab = ba \quad \checkmark$$

Symmetric

$$(a, b)Q(c, d) \quad ad = bc$$

$$(c, d)Q(a, b) \quad cb = da \quad \left. \begin{array}{l} \text{SAME} \\ \checkmark \end{array} \right\}$$

Transitive

$$(a, b)Q(c, d) \quad ad = bc \quad a = \frac{bc}{d}$$

$$(c, d)Q(e, f) \quad cf = de \quad f = \frac{de}{c}$$

$$(a, b)Q(e, f) \quad af = be$$

$$af = \frac{bc}{d} \cdot \frac{de}{c} = be \quad \checkmark$$

Reflexive
Symmetric
& Transitive
so \equiv is
an equivalence
relation!

(b) (2 points) Give three different elements of $[(2, 3)]_Q$.

$$ad = bc$$

$$\frac{a}{b} = \frac{c}{d} \quad \frac{2}{3}, \frac{4}{6}, \frac{8}{12}$$

$(2, 3), (4, 6), (8, 12)$

4. (10 points) Recall that the Fibonacci sequence $\{F_n\}_{n=0}^{\infty}$ is defined by $F_0 = 0, F_1 = 1$, and for $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$. Show that for $n \geq 0$, $\sum_{i=0}^n F_i = F_{n+2} - 1$.

$$F_0 = 0 \quad F_1 = 1 \quad F_n = F_{n-1} + F_{n-2}$$

$$\text{Base case: } \sum_{i=0}^0 F_i = 0 \quad \checkmark$$

$$F_2 - 1 = 1 - 1 = 0$$

$$\sum_{i=0}^1 F_i = 1 \quad \checkmark$$

$$F_3 - 1 = 2 - 1 = 1$$

Inductive step:

$$\text{Assume } \sum_{i=0}^n F_i = F_{n+2} - 1.$$

$$\sum_{i=0}^{n+1} F_i = F_0 + \dots + F_n + F_{n+1}$$

$$= F_{n+2} + F_{n+1} - 1$$

$$\underline{\underline{F_{n+2} + F_{n+1} = F_{n+3}}}$$

$$= F_{n+3} - 1$$

$$= F_{(n+1)+2} - 1 \quad \checkmark$$

$$\text{By induction, } \sum_{i=0}^n F_i = F_{n+2} - 1$$

5. Let X be a set, and let $P = \{(A, B) \in \mathcal{P}(X) \times \mathcal{P}(X) : A \subseteq B\}$ be the set of ordered pairs (A, B) of subsets of X where $A \subseteq B$. Let $\{0, 1, 2\}^X$ denote the set of functions from X to $\{0, 1, 2\}$.

Define a function $F : P \rightarrow \{0, 1, 2\}^X$ by for $(A, B) \in P$ the function $F((A, B)) : X \rightarrow \{0, 1, 2\}$ is defined by for $x \in X$,

$$F((A, B))(x) = \begin{cases} 0 & x \in A \\ 1 & x \in B \text{ and } x \notin A \\ 2 & x \notin B \end{cases}$$

(a) (2 points) Let $X = \{a, b, c, d\}$. What is $F((\{a, d\}, \{a, b, d\}))$? Remember your answer should be a function $X \rightarrow \{0, 1, 2\}$.

$$F((\{a, d\}, \{a, b, d\})) (x) = \begin{cases} 0 & x \in \{a, d\} \\ 1 & x \in \{a, b, d\} \text{ and } x \notin \{a, d\} \\ 2 & x \notin \{a, b, d\} \end{cases}$$

(b) (2 points) Again let $X = \{a, b, c, d\}$. Find a pair (A, B) of subsets of X with $A \subseteq B$ so that $F((A, B))$ is the function $g : X \rightarrow \{0, 1, 2\}$ defined by $g(a) = 1, g(b) = 1, g(c) = 2, g(d) = 2$.

$$A = \{\} \quad B = \{a, b\} \quad \left(\{\}, \{a, b\} \right)$$

Question 5 continued...

- (c) (6 points) Show that for any set X the function $F : P \rightarrow \{0, 1, 2\}^X$ defined on the previous page is a bijection.

surjective (onto)

$$F(A, B)(x) = \begin{cases} 0 & x \in A \\ 1 & x \in B \wedge x \notin A \\ 2 & x \notin B \end{cases}$$

$$0 \rightarrow A$$

$$1 \rightarrow B \cap A^c$$

$$2 \rightarrow (B \cup A)^c$$

These are non overlapping
cases so therefore
for any set X F is surjective

$A^c =$ complement
of A

For any set these three
quantities are different so
it must be surjective

Injective (one to one)

For any set X there is only one ~~set~~
function that it can create, F .

$A, B \cap A^c, (B \cup A)^c$ are non overlapping
so this must be injective!

\therefore injective & surjective

\Rightarrow bijection

This page has been left intentionally blank. You may use it as scratch paper. It will not be graded unless indicated very clearly here and next to the relevant question.

This page has been left intentionally blank. You may use it as scratch paper. It will not be graded unless indicated very clearly here and next to the relevant question.