

# Math 61-1 Midterm 1 version a

EUGENE LO

TOTAL POINTS

**32 / 50**

QUESTION 1

Multiple Choice 10 pts

1.1 2 / 2

- ✓ + 2 pts Correct (b)
- + 0 pts Incorrect

1.2 2 / 2

- ✓ + 2 pts Correct (b)
- + 0 pts Incorrect
- + 2 pts Correct (d)

1.3 0 / 2

- + 2 pts Correct (a)
- ✓ + 0 pts Incorrect

1.4 2 / 2

- ✓ + 2 pts Correct (a)
- + 0 pts Incorrect
- + 2 pts Correct (d)

1.5 0 / 2

- + 2 pts Correct (a)
- ✓ + 0 pts Incorrect

QUESTION 2

Short answer 10 pts

2.1 2 / 2

- ✓ - 0 pts Correct

2.2 0 / 2

- ✓ - 2 pts Incorrect

2.3 0 / 2

- ✓ - 2 pts Incorrect

2.4 0 / 2

- ✓ - 2 pts Incorrect

2.5 2 / 2

- ✓ + 2 pts Correct (5050)
- + 1 pts  $1+2+\dots+100$
- + 1 pts  $n(n+1)/2$
- + 0 pts Incorrect

QUESTION 3

3 Fibonacci numbers 10 / 10

- ✓ - 0 pts Correct

QUESTION 4

Rationals 10 pts

4.1 Equivalence relation 7 / 8

- ✓ - 1 pts Transitivity proof error: how do you know you didn't divide by 0?

☞ Should be " $xy = yx$ " for reflexivity

4.2 Equivalence classes 1.5 / 2

- ✓ - 0.5 pts Claimed  $|[(2,3)]| = 3$

QUESTION 5

Pairs of subsets 10 pts

5.1 Example of the function 2 / 2

- ✓ - 0 pts Correct

5.2 Example of the subsets 1.5 / 2

- ✓ - 0.5 pts the empty set shouldn't have brackets around it

5.3 It is a bijection 0 / 6

- ✓ - 3 pts issue in surjectivity

✓ - **3 pts** issue in injectivity

- You seem to be claiming that the range of  $F$  consists of bijections. This is not true, and it is not what you are trying to show.

# Midterm 1

Last Name: LoFirst Name: EugeneStudent ID: 905108982

Section:            Tuesday:            Thursday:

1A	1B	TA: Albert Zheng
1C	1D	TA: Benjamin Spitz
1E	1F	TA: Eilon Reisin-Tzur

**Instructions:** Do not open this exam until instructed to do so. You will have 50 minutes to complete the exam. Please print your name and student ID number above, and circle the number of your discussion section. **You may not use calculators, books, notes, or any other material to help you. Please make sure your phone is silenced and stowed where you cannot see it. Remember that you are bound by a conduct code, and that you may not look at anyone's paper or let anyone look at your paper, nor may you work on the exam after it has ended. There is some scratch paper at the back of the exam, if you do work there indicate so on the relevant question. Do not write on the backs of the pages. Please circle or box your final answers.**

**Please get out your id and be ready to show it when you turn in your exam.**


Please do not write below this line.

Question	Points	Score
1	10	
2	10	
3	10	
4	10	
5	10	
Total:	50	

1. (10 points) Circle the correct answer (only one answer is correct for each question)


1. If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are functions and  $g \circ f$  is a bijection, then:

- (a)  $f$  must be onto
- (b)  $f$  must be one-to-one and  $g$  must be onto
- (c)  $g$  must be one-to-one
- (d)  $f$  and  $g$  must both be bijections




2. The relation on the integers defined by  $xRy$  if  $x - y$  is odd is:

- (a) reflexive, but not symmetric or transitive
- (b) symmetric, but not transitive or reflexive
- (c) symmetric and transitive, but not reflexive
- (d) an equivalence relation




3. If  $f : X \rightarrow Y$  is an one-to-one function and  $g : Y \rightarrow Z$  is an onto function, then  $g \circ f$  is necessarily:

- (a) not necessarily one-to-one or onto
- (b) one-to-one
- (c) a bijection
- (d) onto




4. For  $X$  a set with  $n$  elements, how many relations are there on  $X$  that are both partial orders and equivalence relations?

- (a) exactly 1
- (b) more than  $n$
- (c) more than 1 and less than  $n$
- (d) exactly  $n$



5. The function  $f : \{a, b\}^* \times \{a, b\}^* \rightarrow \{a, b\}^*$  ( $X^*$  is the set of strings in  $X$ ) defined by  $f((\alpha, \beta)) = \alpha\beta$  is:

- (a) onto but not one-to-one
- (b) one-to-one but not onto
- (c) neither one-to-one nor onto
- (d) onto and one-to-one



2. In this question write down your answer, no need for any justification.


- (a) (2 points) List elements of the equivalence relation on  $\{a, b, c, d, e\}$  determined by the partition  $\{\{a, b\}, \{c\}, \{d, e\}\}$ .

$$R = \{(a, a), (b, b), (a, b), (b, a), (c, c), (d, d), (e, e), (d, e), (e, d)\}$$

- (b) (2 points) Give an example of a relation on  $X = \{a, b, c\}$  that is reflexive and symmetric but not transitive.

$$x, y \in X \quad xRy \text{ if } R = \{(a, a), (b, b), (c, c), (a, b), (b, a)\}$$

- (c) (2 points) Give an example of a functions  $f : X \rightarrow Y, g : Y \rightarrow Z$  so that  $g$  is onto but  $g \circ f$  is not onto.



- (d) (2 points) If  $X$  is a set with  $n$  elements where  $n \geq 1$ , how many onto functions are there from  $X$  to  $\{0, 1\}$ ?

$\{0, 1\}$  2

- (e) (2 points) What is  $\sum_{i=0}^{100} i$ ? Feel free to write your answer as the product or sum of a few numbers.

$$S = \sum_{i=0}^{100} i = 0 + 1 + 2 + \dots + 100 = 0 + \frac{100(100+1)}{2} = 5050$$

$1+2 = \frac{2(2+1)}{2}$   
 $1+2+3 = \frac{3(3+1)}{2}$   
 $\dots$   
 $101 \times 100$   
 $\frac{10100}{2}$

$2 \frac{5050}{10100}$   
 $\frac{10100}{2}$

3. (10 points) Recall that the Fibonacci sequence  $\{F_n\}_{n=0}^{\infty}$  is defined by  $F_0 = 0, F_1 = 1$ , and for  $n \geq 2$ ,  $F_n = F_{n-1} + F_{n-2}$ . Show that for  $n \geq 0$ ,  $\sum_{i=0}^n F_i = F_{n+2} - 1$ .

base case:  $n=0$   $\sum_{i=0}^0 F_i = F_{0+2} - 1$   $F_0 = 0$   
 $F_1 = 1$   
 $F_2 = 1$

$F_0 = F_2 - 1$   
 $F_0 = F_1 + F_0 - 1$   
 $F_0 = 1 + 0 - 1 = 0$   
 $0 = 0 \checkmark$  base case is true

inductive step: Assume  $\sum_{i=0}^n F_i = F_{n+2} - 1$  is true.  
 want to show  $\sum_{i=0}^{n+1} F_i = F_{(n+1)+2} - 1$  is true

$\downarrow$

$\sum_{i=0}^n F_i + F_{n+1} = F_{(n+1)+2} - 1$

$F_{n+2} - 1 + F_{n+1} = F_{(n+1)+2} - 1$

given:  $F_n = F_{n-1} + F_{n-2}$  for  $n \geq 2$   
 because  $n \geq 0$ ,  $(n+1)+2$  must be  $\geq 2$ ,

$\frac{F_{n+2} + F_{n+1} - 1}{\downarrow} = F_{(n+1)+2} - 1$

$F_{(n+1)+2} = F_{n+1} + F_{n+2}$

$F_{(n+1)+2} - 1 = F_{(n+1)+2} - 1$

$\square$  done by induction  $\checkmark$

4. Let  $X = \mathbb{Z} \times \mathbb{Z}_{>0}$  (i.e. ordered pairs of integers where the second integer is positive), and define a relation  $Q$  on  $X$  by  $(a, b)Q(c, d)$  if  $ad = bc$ .

(a) (8 points) Show that  $Q$  is an equivalence relation on  $X$ .

reflexive: For all  $(x, y) \in X$ ,  $(x, y)Q(x, y)$   
 $(x, y)Q(x, y) \rightarrow xy = xy \checkmark$   
 $Q$  is reflexive

symmetric: If  $(a, b)Q(c, d)$ , where  $(a, b), (c, d) \in X$   
 then  $(c, d)Q(a, b)$   
 $ad = bc$   
 $cb = ad \rightarrow ad = bc \checkmark$   
 $Q$  is symmetric

transitive: If  $(a, b)Q(c, d)$  and  $(c, d)Q(e, f)$ ,  
 then  $(a, b)Q(e, f)$   
 $(a, b)Q(c, d) \rightarrow \frac{ad}{c} = bc$   
 $(c, d)Q(e, f) \rightarrow cf = de \rightarrow \frac{cf}{e} = d$   
 $(a, b)Q(e, f) \rightarrow af = be$   
 $a\left(\frac{cf}{e}\right) = bc$   
 $acf = bce$   
 $af = be \checkmark$   
 $Q$  is reflexive, symmetric, & transitive: eq. relation. Thus,  $(a, b)Q(e, f)$   
 $Q$  is

(b) (2 points) Give three different elements of  $[(2, 3)]_Q$ .

$(a, b)Q(c, d) \rightarrow ad = bc$   
 $[(2, 3)]_Q = \{(4, 6), (6, 9), (8, 12)\}$

5. Let  $X$  be a set, and let  $P = \{(A, B) \in \mathcal{P}(X) \times \mathcal{P}(X) : A \subseteq B\}$  be the set of ordered pairs  $(A, B)$  of subsets of  $X$  where  $A \subseteq B$ . Let  $\{0, 1, 2\}^X$  denote the set of functions from  $X$  to  $\{0, 1, 2\}$ .

Define a function  $F : P \rightarrow \{0, 1, 2\}^X$  by for  $(A, B) \in P$  the function  $F((A, B)) : X \rightarrow \{0, 1, 2\}$  is defined by for  $x \in X$ ,

$$F((A, B))(x) = \begin{cases} 0 & x \in A \\ 1 & x \in B \text{ and } x \notin A \\ 2 & x \notin B \end{cases}$$

- (a) (2 points) Let  $X = \{a, b, c, d\}$ . What is  $F((\{a, d\}, \{a, b, d\}))$ ? Remember your answer should be a function  $X \rightarrow \{0, 1, 2\}$ .

$$F((\{a, d\}, \{a, b, d\})) \quad \{a, d\} \subseteq \{a, b, d\}$$

$$F((\{a, d\}, \{a, b, d\}))(a) = 0$$

$$F((\{a, d\}, \{a, b, d\}))(b) = 1$$

$$F((\{a, d\}, \{a, b, d\}))(c) = 2$$

$$F((\{a, d\}, \{a, b, d\}))(d) = 0$$

- (b) (2 points) Again let  $X = \{a, b, c, d\}$ . Find a pair  $(A, B)$  of subsets of  $X$  with  $A \subseteq B$  so that  $F((A, B))$  is the function  $g : X \rightarrow \{0, 1, 2\}$  defined by  $g(a) = 1, g(b) = 1, g(c) = 2, g(d) = 2$ .

$$g(a) = 1, g(b) = 1, g(c) = 2, g(d) = 2$$

$$A = \{\emptyset\}$$

$$B = \{a, b\}$$



Question 5 continued...

(c) (6 points) Show that for any set  $X$  the function  $F: P \rightarrow \{0, 1, 2\}^X$  defined on the previous page is a bijection.

let  $F(A, B)$  be the function  $g: X \rightarrow \{0, 1, 2\}$

$F: P \rightarrow \{0, 1, 2\}^X$  is one-to-one s.t. if

$$g(x_1) = g(x_2), \text{ then } x_1 = x_2$$

$$g(x_1) = F(A, B)(x_1) = \begin{cases} 0 & x_1 \in A \\ 1 & x_1 \in B \text{ and } x_1 \notin A \\ 2 & x_1 \notin B \end{cases}$$

$$g(x_2) = F(A, B)(x_2) = \begin{cases} 0 & x_2 \in A \\ 1 & x_2 \in B \text{ and } x_2 \notin A \\ 2 & x_2 \notin B \end{cases}$$

if  $X = \{a, b, c, d\}$  and  $F(\{a, b\}, \{a, b, d\})$

$x_1$  has to equal  $x_2$  s.t.  $g(x_1) = g(x_2)$

one-to-one ✓

Onto: For all  $y \in \{0, 1, 2\}$ , there exists  $x \in X$  s.t.  $g(x) = y$

$$g(x) = F(A, B)(x) = y$$

$g(x)$  MUST correspond to one of the #'s in  $\{0, 1, 2\}$

if  $X = \{a, b, c, d\}$  and  $F(\{a, b\}, \{a, b, d\})$

$$g(a) = 0, \quad g(b) = 1, \quad g(c) = 2, \quad g(d) = 0$$

each  $x \in X$  corresponds to all elements

of  $\{0, 1, 2\}^X$ . If  $x \in A$ , then  $g(x) = 0$ .

(This also includes empty set, b/c  $\{\emptyset\} \subseteq P(X)$ )

otherwise, if  $x \in B$  and  $x \notin A$  then

$x = 1$ , if  $x \notin B$ , then  $x = 2$ .

Thus?  $F: P \rightarrow \{0, 1, 2\}^X$  is a bijection.

This page has been left intentionally blank. You may use it as scratch paper. It will not be graded unless indicated very clearly here and next to the relevant question.

This page has been left intentionally blank. You may use it as scratch paper. It will not be graded unless indicated very clearly here and next to the relevant question.