Math 61 Midterm 2

Derek Hu

TOTAL POINTS

78 / 100

QUESTION 1

1 8 / 10

- \checkmark + 2 pts a) False
- \checkmark + 2 pts b) False
- \checkmark + 2 pts c) False
- ✓ + 2 pts d) True
 - + 2 pts e) True

QUESTION 2

30 pts

2.1 a 6 / 6

✓ - 0 pts Correct

2.2 b 4 / 4

 \checkmark - **0 pts** Correct (based on answer to a)

2.3 C 0 / 10

✓ - 10 pts No significant progress

2.4 d 10 / 10

- ✓ + 10 pts Correct
 - + 2 pts correct constant on (-1)^n
 - + 2 pts correct constant on 5^n
 - + 6 pts g(n)=-2
 - + **2 pts** g(n)=c
 - 2 pts Minor computational error
 - + **0 pts** No significant progress

QUESTION 3

3 **18 / 20**

\checkmark - 2 pts Didn't update the label on d

QUESTION 4 25 pts

4.1 a 10 / 10

✓ + 10 pts Correct

+ 0 pts Did not draw a Hamiltonian cycle

4.2 b 15 / 15

✓ + 15 pts Correct

+ **0 pts** Solution lacked any significant content/tried to use counting method to conclude that no path exists (and believed it worked/used it improperly)

- 8 pts Solution is vaguely worded, incomplete or includes false claims that doesn't affect final argument.

- **3 pts** Does not explain why certain edges must be used/excluded in a potential Hamiltonian cycle.

QUESTION 5

15 pts

5.1 a 2 / 10

- 8 pts Bin 4: The argument does not show any promise of leading to a proof or has not made enough progress to demonstrate the student could have worked out the remaining details. (Usually this is a poor attempt at pigeonhole principle, e.g. one vertex is connected to 5, the next is connected to 5, so that the total is 10; or, concluded # edges > # vertices but tried to say that implies connected)

5.2 b 5 / 5 √ - 0 pts Correct

MIDTERM 2 (MATH 61) FRIDAY, MAY 17TH

Name: Derek Hu

ID: 805 110 848

Circle your discussion section:

Tuesday	Thursday	
2A	(2B)	TA: Harris Khan
2C	2D	TA: Fred Vu
$2\mathrm{E}$	$2\mathrm{F}$	TA: Matthew Stone

This exam has 8 pages, including the cover page. Please make sure your exam includes each page. Please write your name on *each* page you submit. You will have 50 minutes to complete this exam. You may not use a calculator, or consult your textbook, class notes, or any other materials. If you need scratch paper or more space for your answers, please use the back of the pages.

If there is any work on the backs of the pages which you would like to have graded, please indicate this clearly on the front of the page for the corresponding problem.

Show your work for these problems, don't just give an answer. If a question asks you to prove something, please write a complete proof. Unless otherwise stated, you may use any results proved in class or in the textbook, but please make it clear when you are doing so. Unless otherwise stated, you will *not* receive full credit for giving the correct answer with no explanation. You may still earn partial credit even if your final answer is incorrect.

Remember that you are bound by a conduct code, and that you may not look at anyone's paper or let anyone look at your paper.

Question	Points	Score
1	10	
2	30	
3	20	
4	25	
5	15	
Total:	100	

Name:_____

1. [10 pts, 2 points each] Mark each of the following statements as either TRUE or FALSE. For this question you do not need to show any work beyond the final answer.

Be sure to read the questions carefully!



3



. ZS: (v;) = Zm

. Name:_____

2. [30 pts] Solve the following recursion relations. In each case, your answer should be a formula for the n^{th} term of the sequence in terms of n.

Show your work. You may use results proved in class or in the textbook, but make it clear how you are getting your answers. A correct answer on its own will not be sufficient for full credit.

(a) [6 pts] Find a general solution to the recursion relation $a_n = 4a_{n-1} + 5a_{n-2}$. (That is, find a formula for a_n in terms of n and some unknown constants, that will work for any choice of initial conditions.)

Let
$$a_{n}=r^{n}$$
 $(r^{n}=4r^{n-1}+5r^{n-2})$ r^{n-2}
 $r^{2}-4r-5=0$
 $(r-5)(r+1)=0$
 $r=5$ or $r=1$
Leneral Subultan : $[a_{n}=A(5^{n}+B(c))^{n}]$

(b) [4 pts] Find the solution to the recurrence relation $a_n = 4a_{n-1} + 5a_{n-2}$ with initial conditions $a_0 = 5$ and $a_1 = 7$.

$$A_{0} = A \cdot S^{\circ} + B \cdot (E1)^{\circ} = 5$$

$$a_{1} = A \cdot S' + B (E1)' = 7$$

$$A + B = 5$$

$$f = 5A - B = 7$$

$$GA = 12$$

$$A = 7$$

$$B = 3$$

3

• Name:__

3 - (n-1)

- (c) [10 pts] Find the solution to the recurrence relation $b_n = b_{n-1}^4 b_{n-2}^5$ with initial conditions $b_0 = 1$ and $b_1 = 16$.
 - Let an = bn

$$\begin{aligned} \begin{aligned} & |d| \ c_{n}: r^{n} \\ U_{n}: r^{2} = 4r + S \\ r^{2} - 4r - S = 0 \\ (t-S)(r+1) \ge 0 \\ r^{2} - 3n r^{2} - 1 \\ (t-S)(r+1) \ge 0 \\ r \ge S = n r^{2} - 1 \\ U_{n}: A S^{n} + B (-1)^{n} \\ U_{n}: A S^{n} + B (-1)^{n} - 2 \\ (t-S)(r+1) = 0 \\ r \ge S = n r^{2} - 1 \\ U_{n}: A S^{n} + B (-1)^{n} - 2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{0} = 4D_{0} + 16 \\ -8D_{0} = 16 \\ D_{0} = -2 \\ (t-S)(r+1) = 0 \\ D_{0} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = -2 \\ (t-S)(r+1) = 0 \\ D_{2} = -2 \\ (t-S)(r+1) = 0 \\ D_{1} = -2 \\ (t-S)(r+1) = 0 \\ (t-S)(r+1) = 0 \\ (t-S)(r+1) = 0 \\ (t-S)(r+1) = 0$$

i farmer Martin and a second state and a second second second second second second second second second second

т. , , Name:_

3. [20 pts] Use Dijkstra's algorithm to find the length of the shortest path (i.e. the path for which the sum of the labels is as small as possible) between a and z in the weighted graph below. You do not need to find the shortest path, finding it's length will be sufficient.



Show each step of Dijkstra's algorithm. A correct final answer with no work shown will not be sufficient for full credit. Use the blank graphs below for your answer. If you make a mistake, clearly cross it out and continue using the next blank graph. There are additional blank graphs on the back of this page.





















Name:___

- 4. [25 pts] In each of the following graphs, either find a Hamiltonian cycle (i.e. a cycle which uses every vertex exactly once), or prove that the graph does not have a Hamiltonian cycle.
 - If the graph does have a Hamiltonian cycle, **CLEARLY** drawing this cycle on the provided graph (so that there's no ambiguity as to which edges are used, and in which order), or listing out the vertices in the order traveled (i.e. writing something like (a, b, e, d, c, a)) will be sufficient for full credit.

If the graph does not have a Hamiltonian cycle, you must give an explanation as to why. Simply drawing diagrams with no explanation will NOT be sufficient for full credit.

(a) [10 pts]



a, d, i, l have degree 2 so all fleir edges an included

{a; 5, c, d, h, l, k, g, f, j, i, e, a}

. .

Name:

(b) [15 pts]



Armilfonian cycle. Vertices h & j have No 1 degree of 2 so edges (h,e), (h,i), (ji), and (jig) must be included in the cycle. After including these edges, vertex is has a degree of 2, so (i,e), (i,f), \$ (1,9) cannot be indeided in the cycle. The only 2 possible edges left to be included that to cident @ F an (F,e) & (F,g) (since every vertex in - a humiltonian cycle has exactly 2 edges incident to It), so (f, e) \$ (f,g) ourst be included to create a hamiltonian cycle, it Men is one at all. However, at her (f,e) & (Fig) are included, a cycle is already dealed "chann above) & thus a hamiltonian cycle for the entite graph is not possible



Name:_

5. [15 pts]

(a) [10 pts] Let G be a simple graph with 10 vertices, in which every vertex has degree at least 5. Prove that G is connected. [Hint: You may want to use the pigeonhole principle.]

(b) [5 pts] Draw a *simple* graph with 10 vertices which is *not* connected, in which every vertex has degree 4.



8

