

MIDTERM 1 (MATH 61)  
MONDAY, OCTOBER 21ST

Name: SADHANA VADREVVU

ID: 205095030

Circle your discussion section:

Tuesday    Thursday

2A

2B

TA: Talon Stark

2C

**2D**

TA: Cameron Kissler

2E

2F

TA: Benjamin Spitz

This exam has 5 (double sided) pages, including the cover page, and a blank page at the end. Please make sure your exam includes each page. Please write your name on *each* page you submit. You will have 50 minutes to complete this exam. **You may not use a calculator**, or consult your textbook, class notes, or any other materials. If you need scratch paper or more space for your answers, please use the extra page at the end.

If there is any work on the blank pages which you would like to have graded, please indicate this **CLEARLY** on the page for the corresponding problem.

Show your work for these problems, don't just give an answer. If a question asks you to prove something, please write a complete proof. Unless otherwise stated, you may use any results proved in class or in the textbook, but please make it clear when you are doing so. Unless otherwise stated, you will *not* receive full credit for giving the correct answer with no explanation. You may still earn partial credit even if your final answer is incorrect.

Remember that you are bound by a conduct code, and that you may not look at anyone's paper or let anyone look at your paper.

| Question | Points | Score |
|----------|--------|-------|
| 1        | 10     |       |
| 2        | 35     |       |
| 3        | 20     |       |
| 4        | 15     |       |
| 5        | 20     |       |
| Total:   | 100    |       |

symmetric:  
if  $(a, b) \in R, (b, a) \in R$

Name: Sadhana

1. [10 pts, 2 points each] Mark each of the following statements as either TRUE or FALSE. For this question you do not need to show any work beyond the final answer.

Be sure to read the questions carefully!

|  |       |
|--|-------|
| (a) The sets $\{3, \clubsuit, \{1, 2\}\}$ and $\{\{2, 1, 1\}, 3, \clubsuit, 3\}$ are equal.  | true  |
| (b) There exists a surjective (i.e. onto) function:<br>$f: \mathcal{P}(\{1, 2, 3, 4\}) \rightarrow \{A, B, C, D, E\} \times \{X, Y, Z\}$<br>where $\mathcal{P}(\{1, 2, 3, 4\})$ is the power set of $\{1, 2, 3, 4\}$ . |       |
| (c) If $X = \{1, 2, 3, 4, 5, 6\}$ then any injective (i.e. one-to-one) function $f: X \rightarrow X$ must also be surjective (i.e. onto).  | false |
| (d) The relation $R$ on $\mathbb{Z}$ defined by $xRy$ if $x \neq y$ is antisymmetric.  |       |
| (e) For any positive integers $n$ and $r$ with $r + 1 \leq n, C(n, r) < C(n, r + 1)$ (i.e. if $X$ is an $n$ -element set, there are always more $(r + 1)$ -combinations of $X$ than $r$ -combinations of $X$ ).        | false |

1 1  
2 2  
3 3  
4 4  
5 5  
6 6

antisymmetric:  
if  $(a, b) \in R$

$$= \frac{6 \cdot 5 \cdot 4 \cdot 2}{3 \cdot 2 \cdot 1} = 20$$

$$\frac{\binom{6}{3}}{\binom{6}{4}} = \frac{6}{2! \cdot 4!} = \frac{3 \cdot 2 \cdot 5}{2 \cdot 1} = 15$$

Name: \_\_\_\_\_

2. [35 pts] Compute the following quantities. You may leave your answers in terms of exponents and factorials, but do not leave your final answers in terms of  $P(n, r)$  or  $C(n, r)$  (so  $4^{12} \frac{15!}{3!6!2!}$  would be an acceptable final answer, but  $P(10, 3)C(18, 7)$  would not).

Show your work. It should be clear how you got your answers.

- (a) [10 pts] The number of ways to form a 10 card hand from a standard 52 card deck (containing 13 clubs, 13 diamonds, 13 hearts and 13 spades) consisting of exactly 5 clubs, 3 diamonds, 2 hearts and no spades. (The order of the cards in this hand is irrelevant, only the set of 10 cards picked.)

                                                                     

- (b) [10 pts] The number of 6 letter strings that can be formed from the letters A, B, C, D, E (allowing repeats) which contain at least one A and at least one B. [Hint: It may be easier to count the number of strings which don't satisfy this.]

total number of 6 letter strings:  
 $5^6$

strings w/o 1 A:  $4^6$

strings w/o 1 B:  $4^6$

Name: \_\_\_\_\_

- (c) [15 pts] The number of permutations of the letters BOOKKEEPER (that is, strings of length 10 containing exactly 1 B, 2 O's, 2 K's, 3 E's, 1 P and 1 R) *such that the first E occurs before the first O.*

[So 'PREBOKEEOK' would be one such permutation, but 'BOPKEREKOE' would not.]

BOOKKEEPER

total num!

Name: \_\_\_\_\_

3. [20 pts] In each problem below, the relation  $R$  is *NOT* an equivalence relation. In each problem, identify a specific property of equivalence relations which fails (either reflexivity, symmetry or transitivity), and give a specific example to prove that it fails.

(a) [5 pts]  $X = \mathcal{P}(\{1, 2, 3, 4, 5\})$ ,  $R$  is the relation on  $X$  defined by  $ARB$  is  $A \subseteq B$  (i.e.  $A$  is a subset of  $B$ ).

(b) [7 pts]  $R$  is the relation on  $\mathbb{R}$  (the set of real numbers) defined by  $xRy$  if  $|x - y| < 1$ .

$$|2 - 2| < 1 \quad (2, 2) \in R$$

$$|3 - 3|$$

Name: \_\_\_\_\_

- (c) [8 pts]  $X = \{A \mid A \subseteq \{1, 2, 3, 4, 5, 6\} \text{ and } |A| = 3\}$  is the set of *three element* subsets of  $\{1, 2, 3, 4, 5, 6\}$ .  $R$  is the relation on  $X$  defined by  $ARB$  if  $A \cap B \neq \emptyset$ .

Name: Sadhana Vadrevu

4. [15 pts] Let  $X, Y$  and  $Z$  be sets, and let  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  be functions, and let  $h = g \circ f$  be the composition of  $f$  and  $g$ . (That is,  $h$  is the function from  $X$  to  $Z$  defined by  $h(x) = g(f(x))$ .)

- (a) [7 pts] Prove that if  $f$  and  $g$  are both onto, then  $h$  is onto as well. [Hint: For any  $z \in Z$ , prove that there is some  $x \in X$  with  $h(x) = z$ .]

If  $f$  is onto, then for every  $y \in Y$  there is some  $x \in X$  where  $f(x) = y$ . And, if  $g$  is onto, then for every  $z \in Z$  there is some  $y \in Y$  where  $g(y) = z$ . So,  $f(x)$  contains all the inputs for  $g$ , and  $g(y)$  contains all the values of  $Z$ .  $h(x) = g(f(x))$ . Because  $g$  is onto, every  $z \in Z$  has a  $y \in Y$  with  $g(y) = z$  so every

- (b) [8 pts] Prove that if  $f$  and  $g$  are both one-to-one, then  $h$  is one-to-one as well. [Hint: Show that if  $h(x_1) = h(x_2)$  for some  $x_1, x_2 \in X$  then  $x_1 = x_2$ .]

If  $f$  is one-to-one,  
then if  $f(x_1) = f(x_2)$ ,  $x_1 = x_2$ .  
If  $g$  is also one-to-one,  
then if  $g(x_3) = g(x_4)$ ,  $x_3 = x_4$ .  
 $h(x_1) = h(x_2)$  means  
 $g(f(x_1)) = g(f(x_2))$ . By our definitions,  
of  $g$  being one-to-one, this means  
 $f(x_1) = f(x_2)$ .  $f$  is also one-to-one,  
so  $f(x_1) = f(x_2)$  implies  $x_1 = x_2$ . ✓

Name: \_\_\_\_\_

5. [20 pts] Prove by induction that for any positive integer  $n$ ,

$$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{(n-1)^2} + \frac{1}{n^2} \leq 2 - \frac{1}{n}.$$

base case:  $n=1$

$$\frac{1}{1^2} = 1 \leq 2 - \frac{1}{1} = 2 - 1 = 1 \checkmark$$

inductive step:

assume  $P(n)$  is true:  $\frac{1}{1^2} + \frac{1}{2^2} + \dots + \frac{1}{n^2} \leq 2 - \frac{1}{n}$

then,

$$\frac{1}{1^2} + \frac{1}{2^2} + \dots + \frac{1}{n^2} + \frac{1}{(n+1)^2} \leq 2 - \frac{1}{n} + \left( \frac{1}{(n+1)^2} \right)$$

$$2 - \left( \frac{1}{n} + \frac{1}{n^2 + 2n + 1} \right)$$

$$2 - \left( \frac{n^2 + 2n + 1}{n(n^2 + 2n + 1)} + \frac{n}{n(n^2 + 2n + 1)} \right)$$

$$2 - \left( \frac{n^2 + 3n + 1}{n(n^2 + 2n + 1)} \right)$$

$$2 - \left( \frac{n^2 + 3n + 1}{n(n+1)^2} \right) \leq 2 - \left( \frac{n^2 + 3n + 1}{(n+1)^2} \right)$$

$$= 2 - \left( \frac{(n+1)^2 + n}{(n+1)^2} \right)$$

$$= 2 - 1 - \frac{n}{(n+1)^2}$$

$$= 1 - \frac{n}{(n+1)^2}$$

$$\leq 2 - \frac{n}{(n+1)^2}$$

$$\leq 2 - \frac{n}{n+1}$$

$$(n+1)(n+1) \\ n^2 + n + n + 1$$



Name: \_\_\_\_\_

This page is intentionally blank. You may use it if you need more room for your solutions.

Name: \_\_\_\_\_

This page is intentionally blank. You may use it if you need more room for your solutions.