

Partial solution

Math 61 Midterm 1
OCT 21 2015

50 minutes

Your Name: _____

UCLA ID: _____

SECTION: Cross one box below

Day \ T.A.	John	Zach	Sam
Tuesday	1A	1C	1E
Thursday	1B	1D	1F

Rules: You MUST simplify completely and BOX all answers with an **INK PEN**. You are allowed to use only this paper and pen/pencil. A one-sided hand-written formula sheet is allowed. No calculators, no books, no notebooks, no web access. You MUST write your name and UCLA id. Except for the last problem, you MUST write out your logical reasoning and/or proof in full. You have exactly 50 minutes.

Warning: At 1:50pm you're OUTATIME, those caught writing after time get automatic 10% score deduction.

Problem	Value	Score
Problem 1	8	
Problem 2	10	
Problem 3	10	
Problem 4	12	
Problem 5	10	
Total	50	

Problem 1.

(a) Show that

$$1 \cdot 2^1 + 2 \cdot 2^2 + 3 \cdot 2^3 + \dots + n \cdot 2^n = (n-1) \cdot 2^{n+1} + 2.$$

Induction ① $n=1$ $1 \cdot 2^1 = 2 = (1-1)2^2 + 2 = 2$ ✓

② Assume identity holds for n prove it for $n+1$

i.e. Assume $1 \cdot 2^1 + 2 \cdot 2^2 + \dots + n \cdot 2^n = (n-1) \cdot 2^{n+1} + 2$

show $1 \cdot 2^1 + 2 \cdot 2^2 + \dots + n \cdot 2^n + (n+1) \cdot 2^{n+1} = n \cdot 2^{n+2} + 2$

$$\begin{aligned} 1 \cdot 2^1 + 2 \cdot 2^2 + \dots + n \cdot 2^n + (n+1) \cdot 2^{n+1} &= (1 \cdot 2^1 + \dots + n \cdot 2^n) + (n+1) \cdot 2^{n+1} \\ &= (n-1) \cdot 2^{n+1} + 2 + (n+1) \cdot 2^{n+1} \quad \text{by assumption} \\ &= 2^{n+1} (n-1 + n+1) + 2 \\ &= 2^{n+1} \cdot 2 \cdot n + 2 = 2^{n+2} \cdot n + 2 \end{aligned}$$

by induction identity holds for all n as desired

(b) Show by induction that $e^n \geq n+1$ for integers $n \geq 1$ ($e = 2.71\dots$).

① base case $e^1 = 2.71 \geq 1+1 = 2$ ✓

② assume $e^n \geq n+1$ show $e^{n+1} \geq n+2$

$$e^{n+1} = e^n \cdot e \geq (n+1)e \quad \text{by assumption} \\ = e n + e$$

now $e \cdot n \geq n$ ($e > 1$) and $e > 2$ so $e n + e > n + 2$


so $e^{n+1} > n + 2$

By induction the inequality holds for all n .

Problem 2. Find the number of grid paths from $(0,0)$ to $(6,6)$ that

$(6,6)$

- (a) go through $(2,2)$,
- (b) go through $(4,4)$,
- (c) go through $(2,2)$ or $(4,4)$,
- (d) do not go through $(2,2)$ and $(4,4)$.



You can write your answers in terms of binomials.

$(0,0)$

$$(a) \binom{4}{2} \binom{8}{4}$$

$$(b) \binom{8}{4} \binom{4}{2}$$

$$(c) A = \{\text{paths through } (2,2)\} \quad B = \{\text{paths through } (4,4)\}$$

$$\text{want } |A \cup B| = |A| + |B| - |A \cap B|$$

$$|A| = \binom{4}{2} \binom{8}{4}, \quad |B| = \binom{4}{2} \binom{8}{4}, \quad |A \cap B| = \binom{4}{2} \binom{4}{2} \binom{4}{2}$$

$$|A \cup B| = \binom{4}{2} \binom{8}{4} + \binom{4}{2} \binom{8}{4} - \binom{4}{2}^3$$

$$(d) U = \{\text{all paths}\}, \quad |U| = \binom{12}{6}$$

$$|U - A \cup B| = |U| - |A \cup B| = \binom{12}{6} - \left(2 \binom{4}{2} \binom{8}{4} - \binom{4}{2}^3 \right)$$

Problem 3. Compute the number of 4-subsets of $\{1, 2, 3, \dots, 10\}$ that:

- (a) contain 5,
- (b) contain only one prime number 2, 3, 5, 7,
- (c) the minimum or maximum is 5,
- (d) the product of three of the entries is 6.

You can write your answers in terms of binomials.

$$(a) \binom{9}{3}$$

$$(b) \text{ choice of prime } \times \binom{6}{3} = 4 \cdot \binom{6}{3}$$

$$(c) A = \{S \mid \min(S) = 5\}, \quad B = \{S \mid \max(S) = 5\}$$
$$|A| = \binom{5}{3}, \quad |B| = \binom{4}{3}, \quad |A \cap B| = 0 \text{ since subset has size 4.}$$


$$\text{want } |A \cup B| = |A| + |B| - |A \cap B| = \binom{5}{3} + \binom{4}{3} - 0$$

$$(d) S = \{1, 2, 3, a\} \quad a \in \{4, 5, \dots, 10\}$$


$$\text{so } \binom{7}{1} = 7$$

Problem 4. Let $X = \{(1, 2, 3), (2, 1, 3), (1, 3, 2), (2, 3, 1), (3, 1, 2), (3, 2, 1)\}$ be the set of permutations of size 3. For each of these relations R on X , draw its digraph and decide whether each is reflexive, symmetric or transitive (or neither).


- (a) $(a_1, a_2, a_3)R(b_1, b_2, b_3)$ if and only if $a_1 = b_1$.
- (b) pRq if and only if q can be obtained from p by swapping two adjacent elements.
e.g. $(1, 2, 3)R(2, 1, 3)$.
- (c) The relation induced from the partition $X_1 = \{(1, 2, 3)\}$, $X_2 = \{(2, 1, 3), (3, 2, 1), (1, 3, 2)\}$, $X_3 = \{(2, 3, 1), (3, 1, 2)\}$ of X .



reflexive
symmetric
transitive



not reflexive ex $(1,2,3)$
symmetric
not transitive
 $(1,2,3)R(2,1,3)R(2,3,1)$
but $(1,2,3) \not R (2,3,1)$



reflexive
symmetric
transitive
(R comes from a partition so it is an equivalence relation)

Problem 5. True or False Circle the answers only **with ink**, next to the questions. No reasoning/calculations will be taken into account.

- (a) The sequence $a_n = n! - 2^n$ is decreasing. **T.** or **F.**
- (b) The sequence $1/\binom{2}{2}, 1/\binom{3}{2}, 1/\binom{4}{2}, \dots$ is nonincreasing. **T.** or **F.**
- (c) Given sets $A, B, C \subset U$ the set $A \cup \overline{B \cup C}$ equals the set $(A \cap \overline{B}) \cup (A \cap \overline{C})$. **T.** or **F.**
- (d) The name EMMETT has more than 88 rearrangements of its letters. **T.** or **F.**
- (e) A prime number p divides all the binomial numbers $\binom{p}{1}, \binom{p}{2}, \dots, \binom{p}{p-1}$. **T.** or **F.**
- (f) There are more injections than surjections from $\{A, B, C, D\}$ to $\{1, 2, 3, 4\}$. **T.** or **F.**
- (g) There are more subsets of $\{1, 2, \dots, 11\}$ of odd size than even size. **T.** or **F.**
- (h) There are the same ^{# of} nonnegative integer solutions to $x_1 + x_2 + x_3 = 4$ as positive integer solutions to $y_1 + y_2 + y_3 = 7$. **T.** or **F.**
- (i) The coefficient of x^2y^2 in $(x + y + 1)^6$ is $\binom{6}{4}$. **T.** or **F.**
- (j) There are more symmetric relations than antisymmetric relations on n elements. **T.** or **F.**