

# Physics 1B - Midterm 1

Due Tuesday, April 21 at 11:59 PM on Gradescope

UCLA / Spring 2020 / Brian Naranjo

---

## Problems


### 1) Fluid mechanics

Taking the  $y$  axis to point upward, consider an upright vase of height  $h_0$  whose base is centered on the origin. The vase is rotationally symmetric about the  $y$  axis with radial profile

$$r(y) = \alpha\sqrt{y + y_0},$$

where  $\alpha$  and  $y_0$  are positive constants.

The vase's base is flat with radius  $r(y = 0) = \alpha\sqrt{y_0}$ , and the vase's rim has radius  $r(y = h_0) = \alpha\sqrt{h_0 + y_0}$ . Initially, the vase is filled to the rim with a fluid of density  $\rho$ . The fluid at the top of the vase is exposed to the atmosphere of pressure  $p_0$ . The vase has a small circular drain of radius  $r_0$  (assume  $r_0 \ll \alpha\sqrt{y_0}$  so that the drain is much smaller than the base) at the origin, through which the fluid flows out to the atmosphere, as shown,



In this problem, we will use Bernoulli's principle and the equation of continuity to calculate the time  $\Delta t$  that it takes for all of the fluid to flow out.

- What is the initial total fluid volume  $V$  in the vase?
- In applying Bernoulli's principle to the fluid's top surface and to the vase's drain, we would like to neglect the kinetic energy term evaluated at the fluid's top surface because it is much smaller than the kinetic energy term evaluated at the vase's drain. Confirm this by calculating

$$\frac{[(1/2)\rho v^2]_{\text{top}}}{[(1/2)\rho v^2]_{\text{drain}}},$$

and showing that it is a small value.

Then use Bernoulli's principle to find the initial *volume flow rate*  $Q$  (dimensions of volume per time) out of the drain. If we were able to maintain a volume flow rate at this constant value, how long would it take to drain the vase?

- c) At time  $t$ , the fluid level is at height  $y(t)$ . Derive a relation of the form

$$\frac{dy}{dt} = -f(y).$$

As previously, in your application of Bernoulli's principle, you may drop the kinetic energy term of the fluid's top surface.

- d) The previous differential equation is separable, and may be integrated as

$$\int_0^{\Delta t} dt = \int_0^{h_0} \frac{1}{f(y)} dy.$$


Carry this out to find the total time  $\Delta t$  that it takes for all of the fluid to flow out.

## 2) Periodic motion

A particle of mass  $m$  has potential energy

$$U(x) = U_0 \left[ \frac{3(x - x_0)}{x_0} - \frac{(x - x_1)^3}{x_1^3} \right],$$

where  $U_0$  and  $x_1$  are positive constants, and  $x_0$  is to be determined.



- a) Calculate  $U'(x)$  and  $U''(x)$ . What are the conditions for there to be stable equilibrium at  $x = 0$ ? Find  $x_0$  so that both of these conditions are satisfied.
- b) Use Taylor's theorem to express  $U(x)$ ,

$$U(x) = U(0) + U'(0)x + \frac{1}{2}U''(0)x^2 + \dots$$


up to, and including, the quadratic term. Then, identify the spring constant  $k$  for small oscillations about the origin.

- c) Assuming that the particle feels a viscous damping force of the form  $F = -bv$ , write down the particle's equation of motion expressed in terms of  $\beta \equiv b/(2m)$  and  $\omega_0^2 \equiv k/m$ .

- d) We assume that  $\omega_0 > \beta$ , so that the motion is underdamped. Then, a general solution may be written

$$x(t) = Ae^{-\beta t} \cos(\omega t + \phi),$$

where  $\omega \equiv \sqrt{\omega_0^2 - \beta^2}$ . We are **not** making the light damping approximation in this problem, so that the motion may be heavily damped but still oscillatory. If the initial conditions are  $x(0) = x_0$  and  $x'(0) = 0$ , then solve for  $A$  and  $\tan \phi$ . (Hint:  $A \neq x_0$ ). Possible motion is



### 3) Mechanical waves

Transverse waves on a string are described by a wave function  $y(x, t)$  that consists of sinusoidal left-traveling and right-traveling waves of the same amplitude  $A$ , angular frequency  $\omega$ , and wave number  $k$ ,

$$y(x, t) = A \cos(kx - \omega t) + A \cos(kx + \omega t + \phi)$$

- a) If the string has mass density  $\mu$ , then what is the string's tension  $T$ ?
- b) If there is a **free** boundary condition at  $x = 0$ , then what restriction must  $y(x, t)$  satisfy at  $x = 0$ ? What value of  $\phi$  ensures that this restriction is satisfied? Using this value of  $\phi$  and the trig identity

$$\cos A + \cos B = 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2},$$

factor the wave function into separate spatial and temporal factors, e.g.,

$$y(x, t) = f(x)g(t).$$

- c) In addition to the **free** boundary condition at  $x = 0$ , further assume that there is a **free** boundary condition at  $x = L$ . Only certain values of wave-vector  $k$  are now allowed, which we label  $k_n$  for  $n = 1, 2, 3, \dots$  (here, we only take positive values of  $k_n$ ). Use the boundary condition at  $x = L$  to determine  $k_n$ . Then, use  $v = \omega/k$  to find  $\omega_n$ . What are  $k_n$  and  $\omega_n$ ?
- d) Recall that a standing wave transmits no net power. However, the string definitely does have energy – it just stays in one place. The wave function  $y_n(x, t)$  of the  $n$ th harmonic is obtained by substituting  $k_n$  and  $\omega_n$  back into your expression for  $y(x, t)$  from the end of part b. Calculate the *instantaneous* kinetic energy of the  $n$ th harmonic,

$$K_n(t) = \int_0^L \frac{1}{2}(\mu dx) \left( \frac{\partial y_n}{\partial t} \right)^2.$$