

20W-PHYSICS1B-1 Midterm 2

TOTAL POINTS

92 / 105

QUESTION 1

Problem 1 15 pts

1.1 (A) 0 / 5

+ 5 pts Correct

✓ + 0 pts Incorrect

1.2 (B) 5 / 5

✓ + 5 pts Correct

+ 0 pts Incorrect

1.3 (C) 5 / 5

✓ + 5 pts Correct

+ 0 pts Incorrect

QUESTION 2

Problem 2 30 pts

2.1 (A) 4 / 10

✓ + 2 pts Gauss Law

✓ + 2 pts Correct formula for E

+ 2 pts Correct calculation for E

+ 2 pts Correct formula for Q

+ 2 pts Correct calculation for Q

+ 0 pts Wrong

2.2 (B) 10 / 10

✓ + 2.5 pts Correct formula

✓ + 2.5 pts Correct interpretation as a point charge

✓ + 2.5 pts Correct total charge enclosed for the system

✓ + 2.5 pts Correct calculation

+ 0 pts wrong

2.3 (C) 10 / 10

✓ + 2 pts Correct Initial energy

✓ + 1 pts Correct initial energy calculation

✓ + 2 pts Correct final energy

✓ + 1 pts Correct final energy calculation

✓ + 2 pts Energy conservation equation

✓ + 2 pts Correct calculation

+ 0 pts Wrong

QUESTION 3

Problem 3 30 pts

3.1 (A) 8 / 10

✓ + 2 pts Correct electric field on left

✓ + 2 pts Correct electric field on right

+ 2 pts Correct formula of electric field

✓ + 1 pts Correct field direction between plates

✓ + 2 pts Correct calculation of sigma

✓ + 1 pts Correct calculation of electric field between plates

+ 0 pts Wrong

3.2 (B) 10 / 10

✓ + 3 pts $V = Ed$ or integral

✓ + 4 pts Correct Expression for V

✓ + 3 pts Correct Numerical Value

+ 0 pts Incorrect

3.3 (C) 10 / 10

✓ + 3 pts $E_{[i]} = E_{[f]}$

✓ + 4 pts Correct Energies

✓ + 3 pts Final Answer

+ 0 pts Incorrect

QUESTION 4

Problem 4 30 pts

4.1 (A) 10 / 10

- ✓ + 2 pts Potential Integral
- ✓ + 2 pts Potential Calculations
- ✓ + 2 pts Correct Potential
- ✓ + 2 pts $E = -\text{grad}(V)$
- ✓ + 2 pts Correct E given V
- + 0 pts Incorrect

4.2 (B) 10 / 10

- ✓ + 2 pts Potential Integral
- ✓ + 2 pts Potential Calculations
- ✓ + 2 pts Correct Potential
- ✓ + 2 pts $E = -\text{grad}(V)$
- ✓ + 2 pts Correct E given V
- + 0 pts Incorrect

4.3 (C) 10 / 10

- ✓ + 5 pts E_x is non-zero
- ✓ + 5 pts Reasoning
- + 0 pts Incorrect
- + 3 pts Mention symmetry (only if incorrect answer)

Write your name here:

Write your UCLA ID here

Midterm #2, Physics 1B, Winter 2020

Section 1 - Thomas Dumitrescu

- Please write your name and UID in the boxes on the front page and your name in the boxes at the top of the odd numbered pages.
- Please write your answers within the margins outlined by the boxes on each page.
- if you are using the "additional space" pages, please label them carefully and refer to them within the answer box for the original problem.
- Closed book, **one** 5x3in note card (both sides) allowed.
- Scientific Calculators allowed, no computers or smartphones, please put books and notebooks in your backpacks.
- If a problem is ambiguous, notify the instructor. Clarifications will be written on the blackboard. Check the board occasionally.
- Time for exam: 60 minutes
- There are 4 questions, check that your exam has all 13 pages.
- Useful quantities:

$$\begin{aligned}\epsilon_0 &= 8.85 \times 10^{-12} \text{ C}^2\text{m}^{-2}\text{N}^{-1} \\ m_{\text{electron}} &= 9.11 \times 10^{-31} \text{ kg} \\ m_{\text{proton}} &= 1.67 \times 10^{-27} \text{ kg} \\ q_{\text{electron}} = -q_{\text{proton}} &= -1.602 \times 10^{-19} \text{ C}\end{aligned}$$

Good Luck !!


-additional space for calculation- Please denote exactly which problem you are working on

A large, empty rectangular box with a thin black border, intended for students to show their work on calculations. The box is currently blank.

Write your name here:


Problem 1: [15pts] Concept questions

a) [5pts] A fictitious Gaussian surface (dotted) encloses three identical negative charges. A fourth, positive charge (equal in magnitude to each of the three negative charges inside) is then placed outside the surface as shown. How do the electric flux Φ through the surface and the magnitude $E = |\vec{E}|$ of the electric field at the point P change once the positive charge is introduced?



- They both stay the same.
- Φ increases, but E stays the same.
- Φ decreases, but E stays the same.
- Φ stays the same, but E increases. *
- Φ stays the same, but E decreases.
- Φ and E both increase.
- Φ and E both decrease.

b) [5pts] Consider a regular pentagon, whose corners are labeled 1, 2, 3, 4, 5 (see figure below). A positive charge $+Q$ is placed at every corner. What is the direction of the electric field at the point A, which is exactly halfway between corners 2 and 3.




- (1) Towards corner 2
- (2) Towards corner 3
- (3) Towards corner 4
- (4) Away from corner 4
- (5) Towards corner 5
- (6) Away from corner 5
- (7) The electric field at point A is zero

Write your name here:

[Redacted Name Box]

c) [5pts] The figure below shows equipotential surfaces with their potential given in Volts. If a negative point charge is placed at point P (indicated by the dot on the 0V equipotential) so that it is initially at rest, in which direction would it move?



- Up
- Down
- Left
- Right

(3) Since the charge is placed on the equipotential with $V = 0$, it will remain at rest.

$$E_{\text{total}} = \frac{k Q \left(\frac{r}{R} \right)^3}{r^2}$$

$$dQ = \rho(r) dr$$

$$d(\rho r)^2$$

Problem 2: [30pts] An atomic nucleus can be modeled as a spherically symmetric charge distribution with the following charge density,

$$\rho(r) = \begin{cases} \rho_0 \left(1 - \frac{r}{R}\right) & (r \leq R) \\ 0 & (r > R) \end{cases} \quad \rho_0 = \frac{\rho_0 R}{R} = \frac{\rho_0}{R}$$

Here R is the radius of the nucleus and ρ_0 denotes the charge density at the center of the nucleus (at $r = 0$).

a) [10pts] Find the magnitude and the direction of the electric field everywhere as a function of position. Express your answer in terms of the total charge Q of the nucleus.

$$E = \begin{cases} r \leq R \\ r > R \end{cases}$$

$$k = \frac{1}{4\pi\epsilon_0}$$

$$E_{r > R} = k \frac{Q}{r^2}$$

$$E_{r \leq R} = \frac{k Q r \left(1 - \frac{r}{R}\right)}{R^3}$$

(choose gaussian surface =

$$E_{r \leq R} \cdot 4\pi r^2 = \frac{\rho_0 \left(1 - \frac{r}{R}\right) \cdot \frac{4}{3}\pi r^3}{\epsilon_0}$$

$$E_{r \leq R} = \frac{\rho_0 \left(1 - \frac{r}{R}\right) r}{3 \epsilon_0}$$

Direction: E is always radially outward if $Q > 0$, inward if $Q < 0$

~~Handwritten notes~~

$$\rho(r) \cdot \frac{4}{3}\pi r^3 = Q$$

$$\frac{Q}{\frac{4}{3}\pi r^3} = \rho(r)$$

$$Q_{\text{enc}} (r \leq R) = \rho(r) \cdot \frac{4}{3}\pi r^3$$

$$Q = \int_0^R \rho_0 \left(1 - \frac{r}{R}\right) dr$$

$$E = \int_0^R \frac{k \rho_0 \left(1 - \frac{r}{R}\right)}{\epsilon_0 r^2} dS$$

Write your name here:

b) [10pts] Find the potential difference $V(r) - V(\infty)$ for any point $r > R$ (i.e. outside the nucleus). Explain your reasoning.

Let $V(\infty) = 0$, then $V(r) = \frac{kq}{r}$, as outside the nucleus, its e-field is spherically symmetric and can be treated as a point charge by Gauss' law,

$$\text{so } V(r) - V(\infty) = \frac{q}{4\pi\epsilon_0 \cdot r} \text{ for } r \text{ outside the nucleus.}$$

c) [10pts] A nucleus can decay by emitting an alpha particle, which consists of two protons and two neutrons (i.e. its mass is $4m_{\text{proton}}$ and its charge is $2q_{\text{proton}}$). Quantum physics makes it possible for the alpha particle to jump to a finite distance $r_0 > R$ outside the nucleus. Assuming that the alpha particle starts from rest at $r_0 = 10^{-14}\text{m}$, what is its escape speed far away from the nucleus. Assume that the charge of the nucleus after it has emitted the alpha particle is $Q = 90q_{\text{proton}}$, and that the nucleus is very heavy, so it remains at rest.

$$V_{\text{escape}} \text{ so then } \frac{1}{2} m v_{\text{escape}}^2 = U_{\text{rest}}$$

$$U_I + K_I = U_F + K_F$$

$$U_F = 0, K_I = 0$$


$$U_I = K_F$$

$$v = \sqrt{\frac{2k(90q_{\text{proton}})(2q_{\text{proton}})}{R_0 \cdot 4m_{\text{proton}}}}$$

$$= 3.525 \times 10^7 \text{ m/s}$$

$$\frac{kq_1q_2}{r_0} = \frac{1}{2} (4m_{\text{proton}}) v^2$$

Problem 3: [30pts] Two very thin conducting sheets are parallel disks with radius $r = 0.4\text{m}$ and they are 2cm apart. The left sheet has charge $q = +1.0 \times 10^{-9}\text{C}$, and the right sheet has charge $-q$. You can model the electric fields as if the sheets were of infinite extent.



a) [10pts] What are the electric fields (magnitudes and directions) in the three regions, i.e. to the left of the left sheet, to the right of the right sheet, and in between the sheets?

To the left of the left sheet: $E = 0$

~~Try between: $E = \frac{\sigma}{2\epsilon_0} - \frac{\sigma}{2\epsilon_0} = 0$ (to the right)~~

To the right: $E = 0$

In between: $E = \frac{\sigma}{2\epsilon_0} = \frac{9.947 \times 10^{-10} \text{ C/m}^2}{2\epsilon_0} = \boxed{112.341 \text{ N/C}}$

$\sigma = \frac{q}{A} = \frac{q}{2\pi r^2} = \frac{1 \times 10^{-9} \text{ C}}{2\pi \text{ m}^2} = 9.947 \times 10^{-10} \text{ C/m}^2$

Write your name here:

[Redacted Name]

b) [10pts] What is the potential difference $V_{\text{left}} - V_{\text{right}}$ between the two sheets?

$$V_{\text{left}} - V_{\text{right}} = \frac{\sigma}{\epsilon_0} \cdot d = Ed = 112.344 \cdot 0.02 \text{ m} = \boxed{2.24689 \text{ V}}$$

c) [10pts] An electron e^- is shot through a very small hole in the left sheet with speed v , toward the right. (You can neglect the effect of the hole on the electric fields.) What is the smallest value of v for which the electron manages to reach the negatively charged plate?

$$\Delta U = q \Delta V \quad U_i + K_i = U_f + K_f \quad U_i, K_f = 0$$


If the electron reaches the plate, then $K_i = U_f$ and $\Delta U = \frac{1}{2}mv^2$

$$\frac{1}{2}mv^2 = q \Delta V$$

$$v = \sqrt{\frac{2q \Delta V}{m}} = \sqrt{\frac{2(1.602 \times 10^{-19})(2.247 \text{ V})}{9.11 \times 10^{-31}}}$$

$$= \boxed{888,473 \text{ m/s}}$$

Problem4: [30pts] a) [10pts] Consider a thin ring of charge. The ring has radius R and lies in the xy -plane, centered at the origin (see figure). Assume that the ring has a uniform linear charge density λ . Compute the electrostatic potential $V(z)$ at a point P on the z -axis, as a function of z . Deduce the magnitude and direction of the electric field on the z -axis.



$$dQ = \lambda ds \quad V = k \frac{q}{r}$$

$$E = \int_0^{2\pi R} \frac{k\lambda}{(z^2 + R^2)} \cdot \frac{z}{\sqrt{z^2 + R^2}} ds$$

$$E(z) = \int_0^{2\pi R} \frac{k\lambda z}{(z^2 + R^2)^{3/2}} ds = \frac{2\pi R \lambda \cdot z}{(R^2 + z^2)^{3/2}} \cdot \frac{1}{4\pi \epsilon_0}$$

$$E \text{ field is } = \frac{\lambda R z}{2\pi \epsilon_0 (R^2 + z^2)^{3/2}}$$


away from the ring if $\lambda > 0$, towards if $\lambda < 0$

$$V(z) = \int_0^{2\pi R} \frac{k\lambda ds}{\sqrt{z^2 + R^2}} = \frac{k\lambda}{\sqrt{z^2 + R^2}} \int_0^{2\pi R} ds = \boxed{\frac{2\pi R \lambda}{4\pi \epsilon_0 \sqrt{z^2 + R^2}}}$$

Write your name here:

[Redacted Name Box]

b) [10pts] Now remove the bottom half of the ring in the previous problem, i.e. consider a charged semicircle lying in the upper half of the xy -plane (where $x > 0$, see figure), centered at the origin, with radius R and linear charge density λ . Find the electrostatic potential $V(z)$ at a point P on the z -axis, as a function of z , for this new situation. Use your answer to determine the z -component $E_z(z)$ of the electric field on the z -axis.



$$V(z) = \frac{kQ}{r} = \int_0^{\pi R} \frac{k \lambda ds}{\sqrt{z^2 + R^2}} = \frac{\lambda \pi R}{4\pi \epsilon_0 \sqrt{z^2 + R^2}} = \frac{\lambda R}{4\epsilon_0 \sqrt{z^2 + R^2}}$$

$$E_z(z) = -\frac{dV(z)}{dz} = -\frac{\lambda \pi R}{4\pi \epsilon_0} \frac{d}{dz} \frac{1}{\sqrt{z^2 + R^2}} = \frac{\lambda R z}{4\epsilon_0 (z^2 + R^2)^{3/2}}$$

c) [10pts] Consider the other components $E_x(z)$, $E_y(z)$ at the point P on the z-axis. Explain why one of these components vanishes while the other does not, and which one is which.

E_y vanishes while E_x does not. This is because the half-ring has symmetry about the xz-plane, meaning that the y components from each side cancel out, while it does not have symmetry about the yz-plane, and the x-components do not cancel out.

Write your name here:

-additional space for calculation- Please denote exactly which problem you are working on

z $p_0 (1 - \frac{r}{R})$ or Q

$$E_{r \leq R} = \int_0^R p_0 \frac{r}{R} dr$$
$$= \frac{dp dV}{4\pi \epsilon_0 r^2} = \int_0^R p_0 - \frac{p_0 r}{R}$$
$$= r p_0 = \frac{p_0 r^2}{2R}$$

