


Midterm # 1


1) A rectangular loop of sides $w = 4$ cm and $l = 5$ cm and resistance $R = 10$ Ohm is moved at a constant velocity of $v = 3$ m/s into, through and then out of a uniform $B = 1.25$ T magnetic field as shown in the Figure. The magnetic field region is much wider than the size of the loop.

- a) Find magnitude and direction of the current induced in the circuit as
- 4 i. the circuit is going into the magnetic field counterclockwise $0.15 A \times 10^{-1}$
 - ii. the circuit is totally within the magnetic field but still moving. clockwise $0.15 A$
 - iii. moving out of the field. counterclockwise $0.15 A$
- b) Find magnitude and direction of the magnetic force on the loop as $\vec{l} \times \vec{B}$ $l = 0.04 + 0.05 + 0.04$
- 3 i. the circuit is going into the magnetic field down page $0.03375 N$
 - ii. the circuit is totally within the magnetic field but still moving. up page $0.03375 N$
 - iii. moving out of the field. out of page down $0.03375 N$
- c) Calculate the total energy dissipated in this motion. $(1.5^2 \cdot 10 + 1.5^2 \cdot 10 + 1.5^2 \cdot 10) = 0.675 W$

$\frac{1}{2} R I^2$
 $w \cdot dl$
 $dl = 3$



- 27 2) An electron ($q = -1.6 \times 10^{-19}$ C, $m = 9.11 \times 10^{-31}$ kg) with an instantaneous velocity $v = 1.50 \times 10^6$ j m/s is moving through a region of constant magnetic field directed into the page with $B = -0.25$ k Tesla as shown in figure.



- $F = q\vec{v} \times \vec{B}$
 $\frac{mv^2}{R} = qvB \Rightarrow R = \frac{mv}{qB}$
 $T = \frac{2\pi r}{v}$
- What are the magnitude and direction of the magnetic force acting on the electron? *right ✓*
 - What is the radius of the electron circular trajectory in this magnetic field? *3.4×10^{-5} m 6×10^{-14} N*
 - What is the period of the electron circular motion? *1.13×10^{-10} s 10^{-5} m -2*
 - What is the direction of the electron circular motion (clockwise or anticlockwise) when viewed from above the page? *clockwise ✓*
 - What are the magnitude and direction of the electric field that must be applied if the electron is to move through this region undeflected?

$F = qE$

Electric field points to the right ✓

$$B = \frac{\mu_0 I}{2\pi r}$$

$$B = \frac{\mu_0 I r}{2\pi R^2} \quad ?$$

- 28 3) An electric current is uniformly distributed throughout a long, straight wire that has a diameter of $d = 50$ mm. If the current through the wire is $I_1 = 6.0$ A, calculate
- $r = 0.025$ m
- The magnitude of the magnetic field $r_1 = 20$ mm radially away from the wire center
 - The magnitude of the magnetic field $r_2 = 50$ mm radially away from the wire center
 - What must the current be for this wire to exert an attractive force per unit length of 10^{-3} N/m on another equal wire carrying a current of $I_2 = 10$ A located $r_3 = 100$ mm away from it?

$$a) \quad B = \frac{\mu_0 I r}{2\pi R^2} = \frac{(4\pi \times 10^{-7})(6.0)(.025)}{(2\pi)(.025)^2} = 7.5 \times 10^{-5} \text{ N}$$

$$b) \quad B = \frac{\mu_0 I}{2\pi r} = \frac{(4\pi \times 10^{-7})(6.0)}{(2\pi)(.05)} = 2.4 \times 10^{-5}$$

$$c) \quad 10^{-3} = \frac{F}{L} = IB = \frac{I' \mu_0 I}{2\pi r}$$

