

Physics 1C

Fall 2015

Midterm # 1

30

1) A rectangular loop of sides  $w = 4$  cm and  $l = 5$  cm and resistance  $R = 10$  Ohm is moved at a constant velocity of  $v = 3$  m/s into, through and then out of a uniform  $B = 1.25$  T magnetic field as shown in the Figure. The magnetic field region is much wider than the size of the loop.



- a) Find magnitude and direction of the current induced in the circuit as
- the circuit is going into the magnetic field  $0.015$  A [counterclockwise] 10
  - the circuit is totally within the magnetic field but still moving.  $0$  A
  - moving out of the field.  $0.015$  A [clockwise]
- b) Find magnitude and direction of the magnetic force on the loop as
- the circuit is going into the magnetic field  $7.5 \cdot 10^{-4}$  N [left] 10
  - the circuit is totally within the magnetic field but still moving.  $0$  N
  - moving out of the field.  $7.5 \cdot 10^{-4}$  N [left]
- c) Calculate the total energy dissipated in this motion.  $7.5 \cdot 10^{-5}$  J 10

$\mathcal{E} = \frac{d\Phi}{dt}$   
 $i = \frac{\mathcal{E}}{R}$

$R = 10$

$l = 0.05$

$w = 0.04$



$\mathcal{E} = \frac{d\Phi}{dt}$

a)  $\mathcal{E} = vBw$

$= 3(1.25)(0.04)$

$2.15 = iR$

$i = \frac{2.15}{10} = 0.015$

iii.  $\mathcal{E} = \int \frac{d\Phi}{dt} = \int Bw \frac{dx}{dt} dt$

$\mathcal{E} \cdot t = Bw \cdot l$

$P = \frac{W}{t}$

c.  $W = F \cdot d$

$7.5 \cdot 10^{-4} \cdot (0.05) \cdot 2$

$7.5 \cdot 10^{-5}$

or  $i^2 R \cdot t = 7.5 \cdot 10^{-5}$

b.  $F = iBw$

$= 0.015(1.25)(0.04)$

$7.5 \cdot 10^{-4}$

iii. moving out of same since  $i$  same since  $v$  constant


$\frac{0.05}{v} = t$

$v = 3$

$t = 0.017$

27

2) An electron ( $q = -1.6 \cdot 10^{-19}$  C,  $m = 9.11 \cdot 10^{-31}$  kg) with an instantaneous velocity  $v = 1.50 \cdot 10^6$  j m/s is moving through a region of constant magnetic field directed into the page with  $B = -0.25$  k Tesla as shown in figure.



$v = 1.5 \cdot 10^6$   
 $v = 6.11$   
 $v = \frac{mv}{r}$   
 $v = \frac{mv}{r}$

- What are the magnitude and direction of the magnetic force acting on the electron?
- What is the radius of the electron circular trajectory in this magnetic field?
- What is the period of the electron circular motion?
- What is the direction of the electron circular motion (clockwise or anticlockwise) when viewed from above the page?
- What are the magnitude and direction of the electric field that must be applied if the electron is to move through this region undeflected?

a.  $F = qvB$   
 $= -1.6 \cdot 10^{-19} (1.5 \cdot 10^6) (-0.25)$   
 $= 6 \cdot 10^{-14}$  N [right] ✓

$\frac{mv^2}{r} = qvB$

b.  $\frac{mv^2}{r} = qvB$   
 $r = \frac{mv}{qB} = \frac{9.11 \cdot 10^{-31} (1.5 \cdot 10^6)}{(1.6 \cdot 10^{-19}) (0.25)}$   
 $3.42 \cdot 10^{-5}$  m ✓

c.  $t = \frac{2\pi r}{v}$   
 $t = \frac{2\pi r}{v} = \frac{2\pi (3.42 \cdot 10^{-5})}{1.5 \cdot 10^6}$   
 $1.43 \cdot 10^{-10}$  s ✓

d. clockwise ✓

e.  $F = qE$   
 $6 \cdot 10^{-14} = 1.6 \cdot 10^{-19} E$   
 $E = 3.75 \cdot 10^5$  V/m [left]

-3 wrong direction

303) An electric current is uniformly distributed throughout a long, straight wire that has a diameter of  $d = 50$  mm. If the current through the wire is  $I_1 = 6.0$  A, calculate

- The magnitude of the magnetic field  $r_1 = 20$  mm radially away from the wire center
- The magnitude of the magnetic field  $r_2 = 50$  mm radially away from the wire center
- What must the current be for this wire to exert an attractive force per unit length of  $10^{-3}$  N/m on another equal wire carrying a current of  $I_2 = 10$  A located  $r_3 = 100$  mm away from it?

a. 
$$B = \frac{\mu_0 i r}{2 \pi R^2}$$

$$= \frac{4\pi \cdot 10^{-7} \cdot 6 \cdot (20 \cdot 10^{-3})}{2 \pi \cdot (25 \cdot 10^{-3})^2}$$

$3.84 \cdot 10^{-5} \text{ T}$

b. 
$$\frac{\mu_0 i}{2 \pi r} = \frac{4\pi \cdot 10^{-7} \cdot 6}{2 \pi (50 \cdot 10^{-3})}$$

$2.4 \cdot 10^{-5} \text{ T}$

c. 
$$F = \frac{\mu_0 i_1 i_2}{2 \pi r} \quad F = i_2 B = \frac{i_2 \mu_0 i_1}{2 \pi r}$$

$$10^{-3} = \frac{i_2 (10) \mu_0}{2 \pi (100 \cdot 10^{-3})}$$

$i_2 = 50 \text{ A}$